

ON REPRESENTING LINEAR PROGRAMS BY GRAPH NEURAL NETWORKS

ZIANG CHEN, JIALIN LIU, XINSHANG WANG, JIANFENG LU, AND WOTAO YIN

ABSTRACT. Learning to optimize is a rapidly growing area that aims to solve optimization problems or improve existing optimization algorithms using machine learning (ML). In particular, the graph neural network (GNN) is considered a suitable ML model for optimization problems whose variables and constraints are permutation-invariant, for example, the linear program (LP). While the literature has reported encouraging numerical results, this paper establishes the theoretical foundation of applying GNNs to solving LPs. Given any size limit of LPs, we construct a GNN that maps different LPs to different outputs. We show that properly built GNNs can reliably predict feasibility, boundedness, and an optimal solution for each LP in a broad class. Our proofs are based upon the recently-discovered connections between the Weisfeiler–Lehman isomorphism test and the GNN. To validate our results, we train a simple GNN and present its accuracy in mapping LPs to their feasibilities and solutions.

1. INTRODUCTION

Applying machine learning (ML) techniques to accelerate optimization, also known as *Learning to Optimize (L2O)*, is attracting increasing attention recently. It has been reported in the literature that L2O shows great potentials on both continuous optimization [1, 10, 40] and combinatorial optimization problems [5, 39]. Many of the L2O works train a parameterized model that takes the optimization problem as input and outputs information useful to the classic algorithms, such as a good initial solution and branching decisions [41], and some even directly generate an approximate optimal solution [20].

In these works, one is building an ML model to approximate the mapping from an explicit optimization instance either to its key properties or directly to its solution. The accuracy of this approximation is called the representation power or expressive power of the model. When the approximation is accurate, the model can solve the problem or provide useful information to guide an optimization algorithm. This paper tries to address a fundamental but open theoretical problem for linear programming (LP):

(P0) *Which neural network can represent LP and predict its key properties and solution?*

To clarify, by solution we mean the optimal solution. We hope the answer to this question paves the way toward answering this question for other optimization types.

Date: September 27, 2022.

A major part of the work of Z. Chen was completed during his internship at Alibaba US DAMO Academy. Corresponding author: Jialin Liu, jialin.liu@alibaba-inc.com.

Linear Programming (LP). LP is an important type of optimization problem with a long history and a wide range of applications, such as scheduling [23], signal processing [8], machine learning [13], etc. A general LP problem is defined as:

$$(1.1) \quad \min_{x \in \mathbb{R}^n} c^\top x, \quad \text{s.t. } Ax \circ b, \quad l \leq x \leq u,$$

where $A \in \mathbb{R}^{m \times n}$, $c \in \mathbb{R}^n$, $b \in \mathbb{R}^m$, $l \in (\mathbb{R} \cup \{-\infty\})^n$, $u \in (\mathbb{R} \cup \{+\infty\})^n$, and $\circ \in \{\leq, =, \geq\}^m$. Any LP problems must follow one of the following three cases [7]:

- Infeasible. The feasible set $\mathcal{X}_F := \{x \in \mathbb{R}^n : Ax \circ b, l \leq x \leq u\}$ is empty. In another word, there is no point in \mathbb{R}^n that satisfies the constraints in LP (1.1).
- Unbounded. The feasible set is non-empty, but the objective value can be arbitrarily good, i.e., lower unbounded. For any $R > 0$, there exists an $x \in \mathcal{X}_F$ such that $c^\top x < -R$.
- Feasible and bounded. There exists $x^* \in \mathcal{X}_F$ such that $c^\top x^* \leq c^\top x$ for all $x \in \mathcal{X}_F$. Such x^* is named as an optimal solution, and $c^\top x^*$ is the optimal objective value.

Thus, considering (P0) on the LP problem (1.1), an ideal ML model is expected to be able to predict the three key characteristics of LP: *feasibility, boundedness, and one of its optimal solutions* (if exist), by taking the LP features (A, b, c, l, u, \circ) as input.

Such input has strong mathematical structure. If we swap the position of the i, j -th variable in (1.1), elements in vectors b, c, l, u, \circ and columns of matrix A will be reordered. The reordered features $(\hat{A}, \hat{b}, \hat{c}, \hat{l}, \hat{u}, \hat{\circ})$ actually represent an exact equivalent LP problem with the original one (A, b, c, l, u, \circ) . Such property is named as *permutation invariance*. If we do not explicitly restrict ML models with a permutation invariant structure, the models may overfit the variable/constraint orders of instances in the training set. Motivated by this point, we adopt *Graph Neural Networks (GNNs)* that are permutation invariant naturally.

GNN in L2O. GNN is a type of neural network defined on graphs and widely applied in many areas, for example, recommender systems, traffic, chemistry, etc [51, 53]. Solving optimization problems with GNN is a recent arising topic in the literature [9, 43]. Many graph-related optimization problems, like minimum vertex cover, traveling salesman problem, vehicle routing problem, can be represented and solved approximately with GNN due to their problem structure [16, 26, 29, 32]. Besides that, one may solve a general LP or mixed-integer linear programming (MILP) with the help of GNN. [18] proposed to represent an MILP problem with a bipartite graph, then applied a message-passing GNN¹ on this graph to guide a classic MILP solver. Since that, many approaches have been proposed to guide MILP or LP solvers with GNN [14, 21, 22, 30, 34, 35, 41, 42, 44, 48]. Although encouraging empirical results have been observed, theoretical foundations are still lack for this approach. Specifying (P0), we ask:

$$(P1) \quad \begin{aligned} & \text{Are there GNNs that can predict the feasibility, boundedness,} \\ & \text{and an optimal solution (if exists), for all LPs?} \end{aligned}$$

Related works and contributions. To answer (P1), one needs the theory of *separation power* and *representation power*. Separation power of a neural network (NN) means its ability to distinguish two different inputs. In our settings, a NN with strong separation power means

¹We will abbreviate message-passing GNN as GNN in this paper for simplicity.

that it can outputs different results when it is applied on any two different LPs. Representation power of NN means its ability to approximate functions of interest. A famous example of representation power is the Universal Approximation Theorem [12, 24]: multilayer perceptron (MLP), a specific type of NN, can universally approximate continuous functions on a compact domain. The theory of representation power is established upon the theory of separation power. Only functions with strong enough separation power may possess strong representation power. The power of GNN has been studied in the literature (see [25, 33, 46] for comprehensive surveys), and some theoretical efforts have been made to represent some graph-related optimization problems with GNN [36, 47]. However, there are still gaps to answer question (P1) since the relationships between characteristics of LP and properties of graphs are not well established. Our contributions are listed below:

- (Separation Power). In the literature, it has been shown that the separation power of GNN is equal to the WL test[4, 19, 52]. However, there exist many pairs of LPs that cannot be distinguished by the WL test. We show that those puzzling LP pairs share the same feasibility, boundedness, and even an optimal solution if exists. Thus, GNN has strong enough separation power.
- (Representation Power). To the best of our knowledge, we established the first complete proof that GNN can universally represent a broad class of LPs. More precisely, we prove that, there exist GNNs that can be arbitrarily close to the following three mappings:
 - LP \rightarrow feasibility,
 - LP \rightarrow optimal objective value ($-\infty$ if unbounded and ∞ if infeasible),
 - LP \rightarrow an optimal solution (if exists),
 although they are not continuous functions and cannot be covered by the literature[11, 27, 28, 37, 38].
- (Experimental Validation). We design and conduct experiments that demonstrate the power of GNN on representing LP.

The rest of this paper is organized as follows. In Section 2, we provide preliminaries, including related notions, definitions and concepts. In Section 3, we show our main theoretical results. The sketches of proofs are provided in Section 4. We validate our results with numerical experiments in Section 5 and we conclude this paper with Section 6.

2. PRELIMINARIES

In this section, we present concepts and definitions that will be used throughout this paper. We first describe how to represent an LP with a weighted bipartite graph, then we define GNN on those LP-induced graphs, and finally we further clarify question (P1) with strict mathematical definitions.

2.1. LP represented as weighted bipartite graph. Before representing LPs with graphs, we first define the graph that we will adopt in this paper: *weighted bipartite graph*. A weighted bipartite graph $G = (V \cup W, E)$ consists of a vertex set $V \cup W$ that are divided into two groups V and W with $V \cap W = \emptyset$, and a collection E of weighted edges, where each edge connects

exactly one vertex in V and one vertex in W . Note that there is no edge connecting vertices in the same vertex group. E can also be viewed as a function $E : V \times W \rightarrow \mathbb{R}$. We use $\mathcal{G}_{m,n}$ to denote the collection of all weighted bipartite graphs $G = (V \cup W, E)$ with $|V| = m$ and $|W| = n$. We always write $V = \{v_1, v_2, \dots, v_m\}$, $W = \{w_1, w_2, \dots, w_n\}$, and $E_{i,j} = E(v_i, w_j)$, for $i \in \{1, 2, \dots, m\}$, $j \in \{1, 2, \dots, n\}$.

One can equip each vertex with a feature vector. Throughout this paper, we denote $h_i^V \in \mathcal{H}^V$ as the feature vector of vertex $v_i \in V$ and denote $h_j^W \in \mathcal{H}^W$ as the feature vector of vertex $w_j \in W$, where $\mathcal{H}^V, \mathcal{H}^W$ are feature spaces. Then we define $\mathcal{H}_m^V := (\mathcal{H}^V)^m, \mathcal{H}_n^W := (\mathcal{H}^W)^n$ and concatenate all the vertex features together as $H = (h_1^V, h_2^V, \dots, h_m^V, h_1^W, h_2^W, \dots, h_n^W) \in \mathcal{H}_m^V \times \mathcal{H}_n^W$. Finally, a weighted bipartite graph with vertex features is defined as a tuple $(G, H) \in \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$.

With the concepts described above, one can represent an LP (1.1) as a graph [18]. Each vertex in W represents a variable in LP and each vertex in V represents a constraint. The graph topology is defined with the matrix A in the linear constraint. More specifically, let us set

- Vertex v_i represents the i -th constraint in $Ax \circ b$, and vertex w_j represents the j -th variable x_j .
- Information of constraints is involved in the feature of v_i : $h_i^V = (b_i, \circ_i)$.
- The space of constraint features is defined as $\mathcal{H}^V := \mathbb{R} \times \{\leq, =, \geq\}$.
- Information of variables is involved in the feature of w_j : $h_j^W = (c_j, l_j, u_j)$.
- The space of variable features is defined as $\mathcal{H}^W := \mathbb{R} \times (\mathbb{R} \cup \{-\infty\}) \times (\mathbb{R} \cup \{+\infty\})$.
- The edge connecting v_i and w_j has weight $E_{i,j} = A_{i,j}$.

Then an LP is represented as a graph $(G, H) \in \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$. In the rest of this paper, we coin such graphs as *LP-induced Graphs* or *LP-Graphs* for simplicity. We present an LP instance and its corresponding LP-graph in Figure 1.




FIGURE 1. An example of LP-graph

2.2. Graph neural networks for LP. Now we define GNNs on LP-graphs. The GNNs in this paper always take an LP-Graph as input and the output has two cases:

- The output is a single real number. In this case, GNN is a function $\mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W \rightarrow \mathbb{R}$ and usually used to predict the properties of the whole graph.

- Each vertex in W has an output. Consequently, GNN is a function $\mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W \rightarrow \mathbb{R}^n$. Since W represents variables in LP, GNN is used to predict properties of each variable in this case.

Now we define the GNN structure precisely. First we encode the input features into the embedding space with learnable functions $f_{\text{in}}^V : \mathcal{H}^V \rightarrow \mathbb{R}^{d_0}$ and $f_{\text{in}}^W : \mathcal{H}^W \rightarrow \mathbb{R}^{d_0}$

$$(2.1) \quad h_i^{0,V} = f_{\text{in}}^V(h_i^V), \quad h_j^{0,W} = f_{\text{in}}^W(h_j^W), \quad i = 1, 2, \dots, m, \quad j = 1, 2, \dots, n,$$

where $h_i^{0,V}, h_j^{0,W} \in \mathbb{R}^{d_0}$ are initial embedded vertex features, d_0 is their dimension. Then we choose learnable functions $f_l^V, f_l^W : \mathbb{R}^{d_{l-1}} \rightarrow \mathbb{R}^{d_l}$ and $g_l^V, g_l^W : \mathbb{R}^{d_{l-1}} \times \mathbb{R}^{d_l} \rightarrow \mathbb{R}^{d_l}$ and update the hidden states with²:

$$(2.2) \quad h_i^{l,V} = g_l^V \left(h_i^{l-1,V}, \sum_{j=1}^n E_{i,j} f_l^W(h_j^{l-1,W}) \right), \quad i = 1, 2, \dots, m,$$

$$(2.3) \quad h_j^{l,W} = g_l^W \left(h_j^{l-1,W}, \sum_{i=1}^m E_{i,j} f_l^V(h_i^{l-1,V}) \right), \quad j = 1, 2, \dots, n,$$

where $h_i^{l,V}, h_j^{l,W} \in \mathbb{R}^{d_l}$ are vertex features are layer l ($1 \leq l \leq L$) and their dimensions are d_1, \dots, d_L respectively. The output layer of the single-output GNN is defined with learnable function $f_{\text{out}} : \mathbb{R}^{d_L} \times \mathbb{R}^{d_L} \rightarrow \mathbb{R}$

$$(2.4) \quad y_{\text{out}} = f_{\text{out}} \left(\sum_{i=1}^m h_i^{L,V}, \sum_{j=1}^n h_j^{L,W} \right).$$

The output of the vertex-output GNN is defined with learnable function $f_{\text{out}}^W : \mathbb{R}^{d_L} \times \mathbb{R}^{d_L} \times \mathbb{R}^{d_L} \rightarrow \mathbb{R}$

$$(2.5) \quad y_{\text{out}}(w_j) = f_{\text{out}}^W \left(\sum_{i=1}^m h_i^{L,V}, \sum_{j=1}^n h_j^{L,W}, h_j^{L,W} \right), \quad j = 1, 2, \dots, n.$$

We denote collections of single-output GNNs and vertex-output GNNs with \mathcal{F}_{GNN} and $\mathcal{F}_{\text{GNN}}^W$ respectively.

$$(2.6) \quad \begin{aligned} \mathcal{F}_{\text{GNN}} &= \{F : \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W \rightarrow \mathbb{R} \mid F \text{ yields (2.1), (2.2), (2.3), (2.4)}\}, \\ \mathcal{F}_{\text{GNN}}^W &= \{F : \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W \rightarrow \mathbb{R}^n \mid F \text{ yields (2.1), (2.2), (2.3), (2.5)}\}. \end{aligned}$$

In practice, all the learnable functions in GNN $f_{\text{in}}^V, f_{\text{in}}^W, f_{\text{out}}, f_{\text{out}}^W, \{f_l^V, f_l^W, g_l^V, g_l^W\}_{l=0}^L$ are usually parameterized with multi-linear perceptrons (MLP). In our theoretical analysis, we assume for simplicity that those functions may take all continuous functions on given domains, following the settings in [4, Section C.1]. Thanks to the universal approximation properties of MLP [12, 24], one can extend our theoretical results by taking those learnable functions as large enough MLPs.

²Note that the update rules in (2.2) and (2.3) follow a message-passing way, where each vertex only collects information from its neighbors. Since $E_{i,j} = 0$ if there is no connection between vertices v_i and w_j , the sum operator in (2.2) can be rewritten as $\sum_{j \in \mathcal{N}(v_i)}$, where $\mathcal{N}(v_i)$ denotes the set of neighbors of vertex v_i .

2.3. Revisiting question (P1). With the definitions of LP-Graph and GNN defined above, we revisit question (P1) and give a more precise mathematical description here. First we define three mappings that respectively describe the feasibility, optimal objective value and an optimal solution of an LP.

Feasibility mapping. The feasibility mapping is a classification function

$$(2.7) \quad \Phi_{\text{feas}} : \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W \rightarrow \{0, 1\},$$

where $\Phi_{\text{feas}}(G, H) = 1$ if the LP associated with (G, H) is feasible and $\Phi_{\text{feas}}(G, H) = 0$ otherwise.

Optimal objective value mapping. Denote

$$(2.8) \quad \Phi_{\text{obj}} : \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W \rightarrow \mathbb{R} \cup \{\infty, -\infty\},$$

as the optimal objective value mapping, i.e., for any $(G, H) \in \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$, $\Phi_{\text{obj}}(G, H)$ is the optimal objective value of the LP problem associated with (G, H) .

Remark 2.1. *The optimal objective value of a LP problem can be a real number or $\infty/-\infty$. The “ ∞ ” case corresponds to infeasible problems, while the “ $-\infty$ ” case consists of LP problems whose objective function is unbounded from below in the feasible region. The preimage of all finite real numbers under Φ_{obj} , $\Phi_{\text{obj}}^{-1}(\mathbb{R})$, actually describes all LPs with finite optimal objective value.*

Remark 2.2. *In the case that a LP problem has finite optimal objective value, it is possible that the problem admits multiple optimal solutions. However, the optimal solution with the smallest ℓ_2 -norm must be unique. In fact, if $x \neq x'$ are two different solutions with $\|x\| = \|x'\|$. Then $\frac{1}{2}(x + x')$ is also an optimal solution due to the convexity of LPs, and it holds that $\|\frac{1}{2}(x + x')\|^2 < \frac{1}{2}\|x\|^2 + \frac{1}{2}\|x'\|^2 = \|x\|^2 = \|x'\|^2$, i.e., $\|\frac{1}{2}(x + x')\| < \|x\| = \|x'\|$, where the inequality is strict since $x \neq x'$. Therefore, x and x' cannot be optimal solutions with the smallest ℓ_2 -norm.*

Optimal solution mapping. For any $(G, H) \in \Phi_{\text{obj}}^{-1}(\mathbb{R})$, we have remarked before that the LP problem associated with (G, H) has a unique optimal solution with the smallest ℓ_2 -norm. Let

$$(2.9) \quad \Phi_{\text{solu}} : \Phi_{\text{obj}}^{-1}(\mathbb{R}) \rightarrow \mathbb{R}^n,$$

be the mapping that maps $(G, H) \in \Phi_{\text{obj}}^{-1}(\mathbb{R})$ to the optimal solution with the smallest ℓ_2 -norm.

Invariance and Equivariance. We denote S_m, S_n as the group consisting of all permutations on vertex groups V, W respectively. In another word, S_m involves all permutations on the constraints of LP and S_n involves all permutations on the variables. In this paper, we say a function $F : \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W \rightarrow \mathbb{R}$ is invariant if it satisfies

$$F(G, H) = F((\sigma_V, \sigma_W) * (G, H)), \quad \forall \sigma_V \in S_m, \sigma_W \in S_n,$$

and a function $F_W : \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W \rightarrow \mathbb{R}^n$ is equivariant if it satisfies

$$\sigma_W(F_W(G, H)) = F_W((\sigma_V, \sigma_W) * (G, H)), \quad \forall \sigma_V \in S_m, \sigma_W \in S_n,$$

where $(\sigma_V, \sigma_W) * (G, H)$ is the permuted graph obtained from reordering indices in (G, H) using (σ_V, σ_W) , which is the group action $S_m \times S_n$ on $\mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$. One can check that Φ_{feas} and Φ_{obj} and any $F \in \mathcal{F}_{\text{GNN}}$ are invariant, Φ_{solu} and any $F_W \in \mathcal{F}_{\text{GNN}}^W$ are equivariant.

Question (P1) actually asks: are there GNNs that can accurately approximate Φ_{feas} , Φ_{obj} and Φ_{solu} ? More precisely, does there exist $F \in \mathcal{F}_{\text{GNN}}$ that is arbitrarily close to Φ_{feas} and Φ_{obj} ? And does there exist $F_W \in \mathcal{F}_{\text{GNN}}^W$ that is arbitrarily close to Φ_{solu} ?

3. MAIN RESULTS

This section presents our main theorems that answer question (P1). As we state in the introduction, representation power is built upon separation power in our paper. We first present with the following theorem that GNN has strong enough separation power to represent LP.

Theorem 3.1. *For any two LP problems respectively corresponding to $(G, H), (\hat{G}, \hat{H}) \in \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$, if $F(G, H) = F(\hat{G}, \hat{H}), \forall F \in \mathcal{F}_{\text{GNN}}$, then they share some common characteristics:*

- (i) *Both LP problems are feasible or both are infeasible, i.e., $\Phi_{\text{feas}}(G, H) = \Phi_{\text{feas}}(\hat{G}, \hat{H})$.*
- (ii) *The two LP problems have the same optimal objective value, i.e., $\Phi_{\text{obj}}(G, H) = \Phi_{\text{obj}}(\hat{G}, \hat{H})$.*
- (iii) *If both problems are feasible with finite optimal objective value, then they have the same optimal solution with the smallest ℓ_2 -norm, up to a permutation.*

Furthermore, if $F_W(G, H) = F_W(\hat{G}, \hat{H}), \forall F_W \in \mathcal{F}_{\text{GNN}}^W$, then (iii) holds without taking permutations, i.e., $\Phi_{\text{solu}}(G, H) = \Phi_{\text{solu}}(\hat{G}, \hat{H})$.

This theorem demonstrates that the function spaces \mathcal{F}_{GNN} and $\mathcal{F}_{\text{GNN}}^W$ are rich enough to distinguish the characteristics of LP. Given two LP instances $(G, H), (\hat{G}, \hat{H})$, as long as their feasibility or boundedness are different, there must exist $F \in \mathcal{F}_{\text{GNN}}$ that can distinguish them: $F(G, H) \neq F(\hat{G}, \hat{H})$. Moreover, as long as their optimal solutions with the smallest ℓ_2 -norm are different, there must exist $F_W \in \mathcal{F}_{\text{GNN}}^W$ that can distinguish them: $F_W(G, H) \neq F_W(\hat{G}, \hat{H})$.

With Theorem 3.1 served as a foundation, we can prove that GNN universally approximates the three mappings Φ_{feas} , Φ_{obj} and Φ_{solu} . Before presenting those results, we first define some concepts of the space $\mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$.

Topology and measure. Let us define the topology and the measure on $\mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$, where $\mathcal{H}^V = \mathbb{R} \times \{\leq, =, \geq\}$ and $\mathcal{H}^W = \mathbb{R} \times (\mathbb{R} \cup \{-\infty\}) \times (\mathbb{R} \cup \{+\infty\})$. Throughout this paper, we consider $\mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$ as a topology space with product topology and measurable space with product measure. It's enough to define the topology and measure of each part separately. Since each graph in this paper (without vertex features) can be represented with matrix $A \in \mathbb{R}^{m \times n}$, the graph space $\mathcal{G}_{m,n}$ is isomorphic to the Euclidean space $\mathbb{R}^{m \times n}$: $\mathcal{G}_{m,n} \cong \mathbb{R}^{m \times n}$ and we equip $\mathcal{G}_{m,n}$ with the standard Euclidean topology and the standard Lebesgue measure. The real spaces \mathbb{R} in \mathcal{H}^W and \mathcal{H}^V are also equipped with the standard Euclidean topology and Lebesgue measure. All the discrete spaces $\{\leq, =, \geq\}$, $\{-\infty\}$, and $\{+\infty\}$ have the discrete topology, and all unions are disjoint unions. We equip those spaces with a discrete measure $\mu(S) = |S|$, where $|S|$ is the number of elements in a discrete space S . This finishes the whole definition and we denote $\text{Meas}(\cdot)$ as the measure on $\mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$.

Theorem 3.2. *Let $X \subset \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$ be measurable with finite measure. For any $\epsilon > 0$, there exists some $F \in \mathcal{F}_{GNN}$, such that*

$$\text{Meas}(\{(G, H) \in X : \mathbb{I}_{F(G,H) > 1/2} \neq \Phi_{feas}(G, H)\}) < \epsilon,$$

where \mathbb{I} is the indicator function, i.e., $\mathbb{I}_{F(G,H) > 1/2} = 1$ if $F(G, H) > 1/2$ and $\mathbb{I}_{F(G,H) > 1/2} = 0$ otherwise.

This theorem shows that GNN is a good classifier for LP instances in X as long as X has finite measure. If we use $F(G, H) > 1/2$ as the criteria to predict the feasibility, the classification error rate is controlled by $\epsilon/\text{Meas}(X)$, where ϵ can be arbitrarily small. Furthermore, we show that GNN can perfectly fit any dataset with finite samples, which is presented in the following corollary.

Corollary 3.3. *For any $\mathcal{D} \subset \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$ with finite instances, there exists $F \in \mathcal{F}_{GNN}$ that*

$$\mathbb{I}_{F(G,H) > 1/2} = \Phi_{feas}(G, H), \quad \forall (G, H) \in \mathcal{D}.$$

Besides feasibility, GNN can also approximate the optimal objective value and the optimal solution with the smallest ℓ_2 -norm up to any given precision. The following theorems provide formal statements.

Theorem 3.4. *Let $X \subset \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$ be measurable with finite measure. For any $\epsilon > 0$, there exists $F_1 \in \mathcal{F}_{GNN}$ such that*

$$(3.1) \quad \text{Meas}(\{(G, H) \in X : \mathbb{I}_{F_1(G,H) > 1/2} \neq \mathbb{I}_{\Phi_{obj}(G,H) \in \mathbb{R}}\}) < \epsilon.$$

For any $\epsilon, \delta > 0$, there exists $F_2 \in \mathcal{F}_{GNN}$ such that

$$(3.2) \quad \text{Meas}\left(\{(G, H) \in X \cap \Phi_{obj}^{-1}(\mathbb{R}) : |F_2(G, H) - \Phi_{obj}(G, H)| > \delta\}\right) < \epsilon.$$

Recall the definition of Φ_{obj} in (2.8) that it can take $\{\pm\infty\}$ as its value. Thus, $\Phi_{obj}(G, H) \in \mathbb{R}$ means the LP corresponding to (G, H) is feasible and bounded with a finite optimal objective value, and inequality (3.1) illustrates that GNN can identify those feasible and bounded LPs among the whole set X , up to a given precision ϵ . Inequality (3.2) shows that GNN can also approximate the optimal value. The measure of the set of LP instances of which the optimal value cannot be approximated with δ -precision is controlled by ϵ . The following corollary gives the results on dataset with finite instances.

Corollary 3.5. *Let $\mathcal{D} \subset \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$ be a finite dataset. There exists $F_1 \in \mathcal{F}_{GNN}$ such that*

$$\mathbb{I}_{F_1(G,H) > 1/2} = \mathbb{I}_{\Phi_{obj}(G,H) \in \mathbb{R}}, \quad \forall (G, H) \in \mathcal{D}.$$

For any $\delta > 0$, there exists $F_2 \in \mathcal{F}_{GNN}$, such that

$$|F_2(G, H) - \Phi_{obj}(G, H)| < \delta, \quad \forall (G, H) \in \mathcal{D} \cap \Phi_{obj}^{-1}(\mathbb{R}).$$

Finally, we show that GNN is able to represent the solution mapping Φ_{solu} .

Theorem 3.6. *Let $X \subset \Phi_{obj}^{-1}(\mathbb{R}) \subset \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$ be measurable with finite measure and be closed under actions in $S_m \times S_n$. For any $\epsilon, \delta > 0$, there exists some $F_W \in \mathcal{F}_{GNN}^W$, such that*

$$\text{Meas}(\{(G, H) \in X : \|F(G, H) - \Phi_{solu}(G, H)\| > \delta\}) < \epsilon.$$

Corollary 3.7. *Let $\mathcal{D} \subset \Phi_{obj}^{-1}(\mathbb{R}) \subset \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$ be a **finite dataset**. For any $\delta > 0$, there exists some $F_W \in \mathcal{F}_{GNN}^W$, such that*

$$\|F(G, H) - \Phi_{solu}(G, H)\| < \delta, \quad \forall (G, H) \in \mathcal{D}.$$

4. SKETCH OF PROOF

In this section, we will present a sketch of our proof lines and provide examples to show the intuitions. The full proof lines are presented in the appendix.

Separation power. Separation power of GNN is closely related to the graph isomorphism problem. To identify whether two given graphs are isomorphic, one may apply the Weisfeiler-Lehman (WL) test [50], a widely studied method that can distinguish a broad class of graphs. It has been shown in the literature that the separation power of GNN is actually equal to the standard WL test. Thus, we present some concepts of the WL test before analyzing GNN.

Algorithm 1 The WL test for LP-Graphs (denoted by WL_{LP})

Require: A graph instance $(G, H) \in \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$ and iteration limit $L > 0$.

- 1: Initialize with $C_i^{0,V} = \text{HASH}_{0,V}(h_i^V)$, $C_j^{0,W} = \text{HASH}_{0,W}(h_j^W)$.
 - 2: **for** $l = 1, 2, \dots, L$ **do**
 - 3: $C_i^{l,V} = \text{HASH}_{l,V} \left(C_i^{l-1,V}, \sum_{j=1}^n E_{i,j} \text{HASH}'_{l,W} \left(C_j^{l-1,W} \right) \right)$.
 - 4: $C_j^{l,W} = \text{HASH}_{l,W} \left(C_j^{l-1,W}, \sum_{i=1}^m E_{i,j} \text{HASH}'_{l,V} \left(C_i^{l-1,V} \right) \right)$.
 - 5: **end for**
 - 6: **return** The multisets containing all colors $\{\{C_i^{L,V}\}_{i=0}^m, \{C_j^{L,W}\}_{j=0}^n\}$.
-

To apply the WL test on LP-graphs, we describe a modified WL test in Algorithm 1, that is slightly different with the standard WL test. In Algorithm 1, the hash functions are injective mappings, and multisets are collections of elements that allow multiple appearance of the same element. $\{\{\}\}$ will be used throughout this paper to denote the multisets. We denote Algorithm 1 by $\text{WL}_{LP}(\cdot)$, and we say that two LP-graphs $(G, H), (\hat{G}, \hat{H})$ can be distinguished by Algorithm 1 if and only if there exist a positive integer L and injective hash functions $\{\text{HASH}_{l,V}, \text{HASH}_{l,W}\}_{l=0}^L$ such that

$$\text{WL}_{LP}((G, H), L) \neq \text{WL}_{LP}((\hat{G}, \hat{H}), L).$$

Unfortunately, similar with the standard WL test (see e.g. [15] and references therein), Algorithm 1 does not perfectly solve the graph isomorphism problem on all LP-graphs. There exist infinitely many pairs of non-isomorphic LP-graphs that cannot be distinguished by Algorithm 1. Figure 2 provide such an example.

However, for two LP-graphs that cannot be distinguished by Algorithm 1, they must share some common characteristics even if they are not isomorphic. For example, let us consider the




FIGURE 2. LP-graphs that cannot be distinguished by the WL test since neighbor information of $\{v_i\}$ and $\{w_j\}$ in the two graphs is equal. Based on the graph pair, we construct three pairs of LPs that are both infeasible, both unbounded, both feasible bounded with the same optimal solution, respectively.

6 LP problems in Figure 2. In each of the three columns, the two non-isomorphic LP problems cannot be distinguished by the WL test. It can be checked that the two problems in the same column share some common characteristics. More specifically, both problems in the first column are infeasible; both problems in the second column are feasible but unbounded; both problems in the third column are feasible and bounded with $(1/2, 1/2, 1/2, 1/2)$ as the optimal solution with the smallest ℓ_2 -norm. Actually, this phenomenon does not only happen on the above instances, but also serves as an universal principle for all LP instances. We summarize the results in the following theorem:

Theorem 4.1. *If $(G, H), (\hat{G}, \hat{H}) \in \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$ are not distinguishable by Algorithm 1, then*

$$\Phi_{feas}(G, H) = \Phi_{feas}(\hat{G}, \hat{H}), \quad \Phi_{obj}(G, H) = \Phi_{obj}(\hat{G}, \hat{H}).$$

Furthermore, if $(G, H), (\hat{G}, \hat{H}) \in \Phi_{obj}^{-1}(\mathbb{R})$, it holds that $\Phi_{solu}(G, H) = \sigma_W(\Phi_{solu}(\hat{G}, \hat{H}))$ for some $\sigma_W \in S_n$.

In other words, the above theorem guarantees the sufficient power of the WL test for separating LP problems with different characteristics, including feasibility, optimal objective value, and optimal solution with smallest ℓ_2 -norm (up to permutation). This combined with the following theorem, which states the equivalence of the separation powers of the WL test and GNNs, yields that GNNs also have sufficient separation power for LP-graphs in the above sense.

Theorem 4.2. *For any $(G, H), (\hat{G}, \hat{H}) \in \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$, the followings are equivalent:*

- (i) (G, H) and (\hat{G}, \hat{H}) are not distinguishable by Algorithm 1.
- (ii) $F(G, H) = F(\hat{G}, \hat{H}), \forall F \in \mathcal{F}_{GNN}$.
- (iii) For any $F_W \in \mathcal{F}_{GNN}^W$, there exists $\sigma_W \in S_n$ such that $F_W(G, H) = \sigma_W(F_W(\hat{G}, \hat{H}))$.

Theorem 4.2 extends the results in [4, 19, 52] to the case with the modified WL test (Algorithm 1) and LP-graphs.

Representation power. Based on the separation power of GNNs, one is able to investigate the representation/approximation power of GNNs. To prove Theorems 3.2, 3.4, and 3.6, we first determine the closure of \mathcal{F}_{GNN} or \mathcal{F}_{GNN}^W in the space of invariant/equivariant continuous functions with respect to the sup-norm, which is also named as the *universal approximation*. The result of \mathcal{F}_{GNN} is stated as follows, where $\mathcal{C}(X, \mathbb{R})$ is the collection/algebra of all real-valued continuous function on X . The result of \mathcal{F}_{GNN}^W can be found in the appendix.

Theorem 4.3. *Let $X \subset \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$ be a compact set. For any $\Phi \in \mathcal{C}(X, \mathbb{R})$ that satisfies $\Phi(G, H) = \Phi(\hat{G}, \hat{H})$ for all $(G, H), (\hat{G}, \hat{H})$ that are not distinguishable by Algorithm 1, and for any $\epsilon > 0$, there exists $F \in \mathcal{F}_{GNN}$ such that*

$$\sup_{(G,H) \in X} |\Phi(G, H) - F(G, H)| < \epsilon.$$


Theorem 4.3 can be viewed as an LP-graph version of results in [4, 19]. Roughly speaking, graph neural networks can approximate any invariant continuous function with arbitrarily small error. Although our target mappings $\Phi_{\text{feas}}, \Phi_{\text{obj}}, \Phi_{\text{solu}}$ are not continuous, we prove (in appendix) that they are measurable. Applying Lusin’s theorem [17, Theorem 1.14], we show that GNN can be arbitrarily close to the target mappings except for a small domain.

5. NUMERICAL EXPERIMENTS

We present the numerical results that validate our theoretical results in this section. We generate LP instances with $m = 10$ and $n = 50$ that are possibly infeasible or feasible and bounded. To check whether GNN can predict feasibility, we generate three data sets with 100, 500, 2500 independent LP instances respectively, and call the solver wrapped in `scipy.optimize.linprog` to get the feasibility, optimal value and an optimal solution for each generated LP. To generate enough feasible and bounded LPs to check whether GNN can approximate the optimal objective value and optimal solution, we follow the same approach as before to generate LP randomly and discard those infeasible LPs until the number of LPs reach our requirement. We train GNNs to fit the three LP characteristics by minimizing the distance between GNN-output and those solver-generated answers. The building and the training of the GNNs are implemented using `TensorFlow` and the codes are modified from³ [18]. We set $L = 2$ for all GNNs and those learnable functions $f_{\text{in}}^V, f_{\text{in}}^W, f_{\text{out}}, f_{\text{out}}^W, \{f_l^V, f_l^W, g_l^V, g_l^W\}_{l=0}^L$ are all parameterized with MLPs. Details can be found in the appendix. Our results are reported in Figure 3.

All the errors reported in Figure 3 are training errors since generalization is out of the scope of this paper. In Figure 3a, the “rate of errors” means the proportion of instances with $\mathbb{1}_{F(G,H) > 1/2} \neq \Phi_{\text{feas}}(G, H)$. This metric exactly equals to zeros as long as the number

³Our codes are modified from <https://github.com/ds4dm/learn2branch> and we will release codes.

FIGURE 3. GNN can approximate Φ_{feas} , Φ_{obj} , and Φ_{solu}

of parameters in GNN is large enough, which directly validates Corollary 3.3. With the three curves in Figure 3a combined together, we conclude that such principle does not violate as the number of samples increases. This consists with Theorem 3.2. Mean squared errors in Figures 3b and 3c are respectively defined as $\mathbb{E}_{(G,H)}|F(G,H) - \Phi_{\text{feas}}(G,H)|^2$ and $\mathbb{E}_{(G,H)}\|F(G,H) - \Phi_{\text{feas}}(G,H)\|^2$. Therefore, Figures 3b and 3c validates Theorems 3.4 and 3.6 respectively. Note that all the instances used in Figure 3b are feasible and bounded. Thus, Figure 3b actually only validates (3.2) in Theorem 3.4. However, due to the fact that feasibility of an LP is equal to the boundedness of its dual problem, one may dualize each LP and use the conclusion of Figure 3a to validate (3.1) in Theorem 3.4.

6. CONCLUSIONS

In this work, we show that graph neural networks, as well as the WL test, have sufficient separation power to distinguish linear programming problems with different characteristics. In addition, GNNs can approximate LP feasibility, optimal objective value, and optimal solution with arbitrarily small errors on compact domains or finite datasets. These results guarantee that GNN is a proper class of machine learning models to represent linear programs, and hence contribute to the theoretical foundation in the learning-to-optimize community. Future directions include investing the representation power of graph neural networks for mixed-integer linear programming (MILP), which has been observed with promising experimental results in the literature.

REFERENCES

- [1] Brandon Amos, *Tutorial on amortized optimization for learning to optimize over continuous domains*, arXiv preprint arXiv:2202.00665 (2022).
- [2] Vikraman Arvind, Johannes Köbler, Gaurav Rattan, and Oleg Verbitsky, *On the power of color refinement*, International symposium on fundamentals of computation theory, 2015, pp. 339–350.
- [3] ———, *Graph isomorphism, color refinement, and compactness*, computational complexity **26** (2017), no. 3, 627–685.
- [4] Waiss Azizian and Marc Lelarge, *Expressive power of invariant and equivariant graph neural networks*, International conference on learning representations, 2021.

- [5] Yoshua Bengio, Andrea Lodi, and Antoine Prouvost, *Machine learning for combinatorial optimization: a methodological tour d’horizon*, European Journal of Operational Research **290** (2021), no. 2, 405–421.
- [6] Christoph Berkholz, Paul Bonsma, and Martin Grohe, *Tight lower and upper bounds for the complexity of canonical colour refinement*, Theory of Computing Systems **60** (2017), no. 4, 581–614.
- [7] Dimitris Bertsimas and John N Tsitsiklis, *Introduction to linear optimization*, Vol. 6.
- [8] Emmanuel J Candes and Terence Tao, *Decoding by linear programming*, IEEE transactions on information theory **51** (2005), no. 12, 4203–4215.
- [9] Quentin Cappart, Didier Chételat, Elias B. Khalil, Andrea Lodi, Christopher Morris, and Petar Veličković, *Combinatorial optimization and reasoning with graph neural networks*, Proceedings of the thirtieth international joint conference on artificial intelligence, IJCAI-21, 20218, pp. 4348–4355.
- [10] Tianlong Chen, Xiaohan Chen, Wuyang Chen, Howard Heaton, Jialin Liu, Zhangyang Wang, and Wotao Yin, *Learning to optimize: A primer and a benchmark*, arXiv preprint arXiv:2103.12828 (2021).
- [11] Zhengdao Chen, Soledad Villar, Lei Chen, and Joan Bruna, *On the equivalence between graph isomorphism testing and function approximation with gnns*, Advances in neural information processing systems **32** (2019).
- [12] George Cybenko, *Approximation by superpositions of a sigmoidal function*, Mathematics of control, signals and systems **2** (1989), no. 4, 303–314.
- [13] Antoine Dedieu, Rahul Mazumder, and Haoyue Wang, *Solving l1-regularized svms and related linear programs: Revisiting the effectiveness of column and constraint generation*, Journal of Machine Learning Research **23** (2022), no. 164, 1–41.
- [14] Jian-Ya Ding, Chao Zhang, Lei Shen, Shengyin Li, Bing Wang, Yinghui Xu, and Le Song, *Accelerating primal solution findings for mixed integer programs based on solution prediction*, Proceedings of the aaai conference on artificial intelligence, 2020, pp. 1452–1459.
- [15] Brendan L Douglas, *The weisfeiler-lehman method and graph isomorphism testing*, arXiv preprint arXiv:1101.5211 (2011).
- [16] Iddo Drori, Anant Kharkar, William R Sickinger, Brandon Kates, Qiang Ma, Suwen Ge, Eden Dolev, Brenda Dietrich, David P Williamson, and Madeleine Udell, *Learning to solve combinatorial optimization problems on real-world graphs in linear time*, 2020 19th IEEE International Conference on Machine Learning and Applications (ICMLA), 2020, pp. 19–24.
- [17] Lawrence C Evans and Ronald F Garzepy, *Measure theory and fine properties of functions*, Routledge, 2018.
- [18] Maxime Gasse, Didier Chételat, Nicola Ferroni, Laurent Charlin, and Andrea Lodi, *Exact combinatorial optimization with graph convolutional neural networks*, Advances in Neural Information Processing Systems **32** (2019).
- [19] Floris Geerts and Juan L Reutter, *Expressiveness and approximation properties of graph neural networks*, International conference on learning representations, 2022.
- [20] Karol Gregor and Yann LeCun, *Learning fast approximations of sparse coding*, Proceedings of the 27th international conference on international conference on machine learning, 2010, pp. 399–406.
- [21] Prateek Gupta, Maxime Gasse, Elias Khalil, Pawan Mudigonda, Andrea Lodi, and Yoshua Bengio, *Hybrid models for learning to branch*, Advances in neural information processing systems **33** (2020), 18087–18097.
- [22] Prateek Gupta, Elias B Khalil, Didier Chételat, Maxime Gasse, Yoshua Bengio, Andrea Lodi, and M Pawan Kumar, *Lookback for learning to branch*, arXiv preprint arXiv:2206.14987 (2022).
- [23] Fred Hanssman and Sidney W Hess, *A linear programming approach to production and employment scheduling*, Management science **1** (1960), 46–51.
- [24] Kurt Hornik, Maxwell Stinchcombe, and Halbert White, *Multilayer feedforward networks are universal approximators*, Neural networks **2** (1989), no. 5, 359–366.
- [25] Stefanie Jegelka, *Theory of graph neural networks: Representation and learning*, arXiv preprint arXiv:2204.07697 (2022).

- [26] Chaitanya K Joshi, Thomas Laurent, and Xavier Bresson, *An efficient graph convolutional network technique for the travelling salesman problem*, arXiv preprint arXiv:1906.01227 (2019).
- [27] Nicolas Keriven, Alberto Bietti, and Samuel Vaiter, *On the universality of graph neural networks on large random graphs*, Advances in Neural Information Processing Systems **34** (2021), 6960–6971.
- [28] Nicolas Keriven and Gabriel Peyré, *Universal invariant and equivariant graph neural networks*, Advances in Neural Information Processing Systems **32** (2019).
- [29] Elias Khalil, Hanjun Dai, Yuyu Zhang, Bistra Dilkina, and Le Song, *Learning combinatorial optimization algorithms over graphs*, Advances in neural information processing systems **30** (2017).
- [30] Elias B Khalil, Christopher Morris, and Andrea Lodi, *Mip-gnn: A data-driven framework for guiding combinatorial solvers*, Update **2** (2022), x3.
- [31] Diederik P Kingma and Jimmy Ba, *Adam: A method for stochastic optimization*, arXiv preprint arXiv:1412.6980 (2014).
- [32] Wouter Kool, Herke van Hoof, and Max Welling, *Attention, learn to solve routing problems!*, International conference on learning representations, 2019.
- [33] Pan Li and Jure Leskovec, *The expressive power of graph neural networks*, Graph neural networks: Foundations, frontiers, and applications, 2022, pp. 63–98.
- [34] Xijun Li, Qingyu Qu, Fangzhou Zhu, Jia Zeng, Mingxuan Yuan, Kun Mao, and Jie Wang, *Learning to reformulate for linear programming*, arXiv preprint arXiv:2201.06216 (2022).
- [35] Defeng Liu, Matteo Fischetti, and Andrea Lodi, *Learning to search in local branching*, Proceedings of the aaai conference on artificial intelligence, 2022, pp. 3796–3803.
- [36] Andreas Loukas, *What graph neural networks cannot learn: depth vs width*, International conference on learning representations, 2020.
- [37] Haggai Maron, Heli Ben-Hamu, Hadar Serviansky, and Yaron Lipman, *Provably powerful graph networks*, Advances in neural information processing systems **32** (2019).
- [38] Haggai Maron, Ethan Fetaya, Nimrod Segol, and Yaron Lipman, *On the universality of invariant networks*, International conference on machine learning, 2019, pp. 4363–4371.
- [39] Nina Mazyavkina, Sergey Sviridov, Sergei Ivanov, and Evgeny Burnaev, *Reinforcement learning for combinatorial optimization: A survey*, Computers & Operations Research **134** (2021), 105400.
- [40] Vishal Monga, Yuelong Li, and Yonina C Eldar, *Algorithm unrolling: Interpretable, efficient deep learning for signal and image processing*, IEEE Signal Processing Magazine **38** (2021), no. 2, 18–44.
- [41] Vinod Nair, Sergey Bartunov, Felix Gimeno, Ingrid von Glehn, Pawel Lichocki, Ivan Lobov, Brendan O’Donoghue, Nicolas Sonnerat, Christian Tjandraatmadja, Pengming Wang, Ravichandra Addanki, Tharindi Hapuarachchi, Thomas Keck, James Keeling, Pushmeet Kohli, Ira Ktena, Yujia Li, Oriol Vinyals, and Yori Zwols, *Solving mixed integer programs using neural networks*, arXiv preprint arXiv:2012.13349 (2020).
- [42] Max B Paulus, Giulia Zarpellon, Andreas Krause, Laurent Charlin, and Chris Maddison, *Learning to cut by looking ahead: Cutting plane selection via imitation learning*, International conference on machine learning, 2022, pp. 17584–17600.
- [43] Yun Peng, Byron Choi, and Jianliang Xu, *Graph learning for combinatorial optimization: a survey of state-of-the-art*, Data Science and Engineering **6** (2021), no. 2, 119–141.
- [44] Qingyu Qu, Xijun Li, Yunfan Zhou, Jia Zeng, Mingxuan Yuan, Jie Wang, Jinhu Lv, Kexin Liu, and Kun Mao, *An improved reinforcement learning algorithm for learning to branch*, arXiv preprint arXiv:2201.06213 (2022).
- [45] Walter Rudin, *Functional analysis*, 2nd ed., International series in pure and applied mathematics, McGraw-Hill, 1991.
- [46] Ryoma Sato, *A survey on the expressive power of graph neural networks*, arXiv preprint arXiv:2003.04078 (2020).
- [47] Ryoma Sato, Makoto Yamada, and Hisashi Kashima, *Approximation ratios of graph neural networks for combinatorial problems*, Advances in Neural Information Processing Systems **32** (2019).

- [48] Yunzhuang Shen, Yuan Sun, Andrew Eberhard, and Xiaodong Li, *Learning primal heuristics for mixed integer programs*, 2021 international joint conference on neural networks (ijcnn), 2021, pp. 1–8.
- [49] Vlad Timofte, *Stone–weierstrass theorems revisited*, Journal of Approximation Theory **136** (2005), no. 1, 45–59.
- [50] Boris Weisfeiler and Andrei Leman, *The reduction of a graph to canonical form and the algebra which appears therein*, NTI, Series **2** (1968), no. 9, 12–16.
- [51] Zonghan Wu, Shirui Pan, Fengwen Chen, Guodong Long, Chengqi Zhang, and S Yu Philip, *A comprehensive survey on graph neural networks*, IEEE transactions on neural networks and learning systems **32** (2020), no. 1, 4–24.
- [52] Keyulu Xu, Weihua Hu, Jure Leskovec, and Stefanie Jegelka, *How powerful are graph neural networks?*, International conference on learning representations, 2019.
- [53] Jie Zhou, Ganqu Cui, Shengding Hu, Zhengyan Zhang, Cheng Yang, Zhiyuan Liu, Lifeng Wang, Changcheng Li, and Maosong Sun, *Graph neural networks: A review of methods and applications*, AI Open **1** (2020), 57–81.

APPENDIX A. WEISFEILER-LEHMAN (WL) TEST AND COLOR REFINEMENT

The WL test can be viewed as a coloring refinement procedure if there are no collisions of hash functions and their weighted averages. More specifically, each vertex is colored initially according to the group it belongs to and its feature – two vertices have the same color if and only if they are in the same vertex group and have the same feature. The initial colors are denoted as $C_1^{0,V}, C_2^{0,V}, \dots, C_m^{0,V}, C_1^{0,W}, C_2^{0,W}, \dots, C_n^{0,W}$. Then at iteration l , the set of vertices with the same color at iteration $l-1$ are further partitioned into several subsets according to the colors of their neighbours – two vertices v_i and $v_{i'}$ are in the same subset if and only if $C_i^{l-1,V} = C_{i'}^{l-1,V}$ and for any $C \in \{C_j^{l-1,W} : 1 \leq j \leq n\}$,

$$\sum_{C_j^{l-1,W}=C} E_{i,j} = \sum_{C_j^{l-1,W}=C} E_{i',j},$$

and it is similar for vertices w_j and $w_{j'}$. After such partition/refinement, vertices are associated with the same color if and only if they are in the same subset, which is the coloring at iteration l . This procedure is terminated if the refinement is trivial, meaning that no sets with the same color are partitioned into at least two subsets, i.e., the coloring is stable. For more information about color refinement, we refer to [2, 3, 6].

We then discuss the stable coloring that Algorithm 1 will converge to, for which we made the following definition, where $\mathcal{S} = \{S_1, S_2, \dots, S_s\}$ is called a partition of a set S if $S_1 \cup S_2 \cup \dots \cup S_s = S$ and $S_i \cap S_{i'} = \emptyset, \forall 1 \leq i < i' \leq s$.

Definition A.1 (Stable Partition Pair of Vertices). *Let $G = (V \cup W, E)$ be a weighted bipartite graph with $V = \{v_1, v_2, \dots, v_n\}$, $W = \{w_1, w_2, \dots, w_n\}$, and vertex features $H = (h_1^V, h_2^V, \dots, h_m^V, h_1^W, h_2^W, \dots, h_n^W)$, and let $\mathcal{I} = \{I_1, I_2, \dots, I_s\}$ and $\mathcal{J} = \{J_1, J_2, \dots, J_t\}$ be partitions of $\{1, 2, \dots, m\}$ and $\{1, 2, \dots, n\}$, respectively. We say that $(\mathcal{I}, \mathcal{J})$ is a stable partition pair of vertices for the graph G if the followings are satisfied:*

- (i) $h_i^V = h_{i'}^V$ holds if $i, i' \in I_p$ for some $p \in \{1, 2, \dots, s\}$.
- (ii) $h_j^W = h_{j'}^W$ holds if $j, j' \in J_q$ for some $q \in \{1, 2, \dots, t\}$.
- (iii) For any $p \in \{1, 2, \dots, s\}$, $q \in \{1, 2, \dots, t\}$, and $i, i' \in I_p$, $\sum_{j \in J_q} E_{i,j} = \sum_{j \in J_q} E_{i',j}$.

(iv) For any $p \in \{1, 2, \dots, s\}$, $q \in \{1, 2, \dots, t\}$, and $j, j' \in J_q$, $\sum_{i \in I_p} E_{i,j} = \sum_{i \in I_p} E_{i,j'}$.

We denote $(\mathcal{I}^l, \mathcal{J}^l)$ as the partition pair corresponding the coloring at iteration l of Algorithm 1. Suppose that there are no collisions. Then it is clear that $(\mathcal{I}^{l+1}, \mathcal{J}^{l+1})$ is finer than $(\mathcal{I}^l, \mathcal{J}^l)$, denoted as $(\mathcal{I}^{l+1}, \mathcal{J}^{l+1}) \preceq (\mathcal{I}^l, \mathcal{J}^l)$, which means that for any $I \in \mathcal{I}^{l+1}$ and any $J \in \mathcal{J}^{l+1}$, there exist $I' \in \mathcal{I}^l$ and $J' \in \mathcal{J}^l$ such that $I \subset I'$ and $J \subset J'$. In addition, $(\mathcal{I}^{l+1}, \mathcal{J}^{l+1}) = (\mathcal{I}^l, \mathcal{J}^l)$ if and only if $(\mathcal{I}^l, \mathcal{J}^l)$ is a stable partition pair of vertices. Note that there is at most $O(|V| + |W|)$ iterations leading to strictly finer partition pair, i.e., $(\mathcal{I}^{l+1}, \mathcal{J}^{l+1}) \preceq (\mathcal{I}^l, \mathcal{J}^l)$ but $(\mathcal{I}^{l+1}, \mathcal{J}^{l+1}) \neq (\mathcal{I}^l, \mathcal{J}^l)$. We can immediately obtain the following result:

Theorem A.2. *If there are no collision of hash functions and their weighted averages, then Algorithm 1 terminates at a stable partition pair of vertices in $O(|V| + |W|)$ iterations.*

Furthermore, for every $(G, H) \in \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$, the coarsest stable partition pair of vertices exists and is unique, which can be proved using techniques similar to the proof of [6, Proposition 3]. Algorithm 1 terminates at the unique coarsest stable partition pair. This is because that the coloring in each iteration of Algorithm 1 is always coarser than the unique coarsest stable partition pair (see [6, Proposition 2]).

APPENDIX B. THE SEPARATION POWER OF THE WL TEST

This section gives the proof and some corollaries of Theorem 4.1. First we present some definitions and lemmas, from which the proof of Theorem 4.1 can be immediately derived.

Definition B.1. *Given $(G, H), (\hat{G}, \hat{H}) \in \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$, we say that (G, H) and (\hat{G}, \hat{H}) can be distinguished by the WL test if there exists some $L \in \mathbb{N}$ and some choices of hash functions, $HASH_{0,V}, HASH_{0,W}, HASH_{l,V}, HASH_{l,W}, HASH'_{l,V}$, and $HASH'_{l,W}$, for $l = 1, 2, \dots, L$, such that the multisets of colors at the L -th iteration of the WL test are different for (G, H) and (\hat{G}, \hat{H}) . Let \sim be an equivalence relationship on $\mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$ defined via: $(G, H) \sim (\hat{G}, \hat{H})$ if and only if they can not be distinguished by the WL test.*

It is clear that $(G, H) \sim (\hat{G}, \hat{H})$ if they are isometric, i.e., there exist two permutations, $\sigma_V : \{1, 2, \dots, m\} \rightarrow \{1, 2, \dots, m\}$ and $\sigma_W : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$, such that $E_{\sigma_V(i), \sigma_W(j)} = \hat{E}_{i,j}$, $h_{\sigma_V(i)}^V = \hat{h}_i^V$, and $h_{\sigma_W(j)}^W = \hat{h}_j^W$, for any $i \in \{1, 2, \dots, m\}$ and $j \in \{1, 2, \dots, n\}$. However, not every pair of WL-indistinguishable graphs consists of isometric ones; see Figure 2 for an example. However, for LP problems that cannot be distinguished by the WL test will share some common properties, even if their associated graphs are not isomorphic.

Lemma B.2. *If two weighted bipartite graphs with vertex features corresponding to two LP problems are indistinguishable by the WL test, then either both problems are feasible or both are infeasible. In other words, the WL test can distinguish two LP problems if one of them is feasible while the other one is infeasible.*

Proof of Lemma B.2. Let us consider two LP problems:

$$(B.1) \quad \begin{aligned} \min_{x \in \mathbb{R}^n} \quad & c^\top x, \\ \text{s.t.} \quad & Ax \circ b, \quad l \leq x \leq u, \end{aligned}$$

and

$$(B.2) \quad \begin{aligned} \min_{x \in \mathbb{R}^n} \quad & \hat{c}^\top x, \\ \text{s.t.} \quad & \hat{A}x \hat{\circ} \hat{b}, \quad \hat{l} \leq x \leq \hat{u}. \end{aligned}$$

Let (G, H) and (\hat{G}, \hat{H}) , where $G = (V \cup W, E)$ and $\hat{G} = (V \cup W, \hat{E})$, be the weighted bipartite graphs with vertex features corresponding to (B.1) and (B.2), respectively.

Since $(G, H) \sim (\hat{G}, \hat{H})$, for some choice of hash functions with no collision during Algorithm 1 for (G, H) and (\hat{G}, \hat{H}) , Theorem A.2 guarantees that Algorithm 1 outputs the same stable coloring for (G, H) and (\hat{G}, \hat{H}) up to permutation. More specifically, after doing some permutation, there exist $\mathcal{I} = \{I_1, I_2, \dots, I_s\}$ and $\mathcal{J} = \{J_1, J_2, \dots, J_t\}$ that are partitions of $\{1, 2, \dots, m\}$ and $\{1, 2, \dots, n\}$, respectively, such that the followings hold:

- $(b_i, \circ_i) = (\hat{b}_i, \hat{\circ}_i)$ and is independent of $i \in I_p$, for any $p \in \{1, 2, \dots, s\}$.
- $(l_j, u_j) = (\hat{l}_j, \hat{u}_j)$ and is independent of $j \in J_q$ for any $q \in \{1, 2, \dots, t\}$.
- For any $p \in \{1, 2, \dots, s\}$ and $q \in \{1, 2, \dots, t\}$, $\sum_{j \in J_q} A_{i,j} = \sum_{j \in J_q} \hat{A}_{i,j}$ and is independent of $i \in I_p$.
- For any $p \in \{1, 2, \dots, s\}$ and $q \in \{1, 2, \dots, t\}$, $\sum_{i \in I_p} A_{i,j} = \sum_{i \in I_p} \hat{A}_{i,j}$ and is independent of $j \in J_q$.

Suppose that the problem (B.1) is feasible with $x \in \mathbb{R}^n$ be some point in the feasible region. Define $y \in \mathbb{R}^t$ via $y_q = \frac{1}{|J_q|} \sum_{j \in J_q} x_j$ and $\hat{x} \in \mathbb{R}^n$ via $\hat{x}_j = y_q$, $j \in J_q$. Fix any $p \in \{1, 2, \dots, s\}$ and some $i_0 \in I_p$. It holds for any $i \in I_p$ that

$$\sum_{j=1}^n A_{i,j} x_j \circ_i b_i, \quad \text{i.e.,} \quad \sum_{q=1}^t \sum_{j \in J_q} A_{i,j} x_j \circ_{i_0} b_{i_0},$$

which implies that

$$\frac{1}{|I_p|} \sum_{i \in I_p} \sum_{q=1}^t \sum_{j \in J_q} A_{i,j} x_j = \frac{1}{|I_p|} \sum_{q=1}^t \sum_{j \in J_q} \left(\sum_{i \in I_p} A_{i,j} \right) x_j \circ_{i_0} b_{i_0}.$$

Notice that $\sum_{i \in I_p} A_{i,j}$ is constant for $j \in J_q$, $\forall q \in \{1, 2, \dots, t\}$. Let us denote $\alpha_q = \sum_{i \in I_p} A_{i,j} = \sum_{i \in I_p} \hat{A}_{i,j}$ for any $j \in J_q$. Then it holds that

$$\frac{1}{|I_p|} \sum_{q=1}^t \sum_{j \in J_q} \alpha_q x_j = \frac{1}{|I_p|} \sum_{q=1}^t \sum_{j \in J_q} \alpha_q y_q = \frac{1}{|I_p|} \sum_{q=1}^t \sum_{j \in J_q} \left(\sum_{i \in I_p} \hat{A}_{i,j} \right) y_q \circ_{i_0} b_{i_0}.$$

Note that

$$\frac{1}{|I_p|} \sum_{q=1}^t \sum_{j \in J_q} \left(\sum_{i \in I_p} \hat{A}_{i,j} \right) y_q = \frac{1}{|I_p|} \sum_{i \in I_p} \sum_{q=1}^t \left(\sum_{j \in J_q} \hat{A}_{i,j} \right) y_q,$$

and that $\sum_{j \in J_q} \hat{A}_{i,j}$ is constant for $i \in I_p$. So one can conclude that

$$\sum_{j=1}^n \hat{A}_{i,j} \hat{x}_j = \sum_{q=1}^t \sum_{j \in J_q} \hat{A}_{i,j} \hat{x}_j = \sum_{q=1}^t \left(\sum_{j \in J_q} \hat{A}_{i,j} \right) y_q \circ_i b_i, \quad \forall i \in I_p,$$

which leads to $\hat{A} \hat{x} \hat{o} \hat{b}$. It can also be seen that $\hat{l} = l \leq \hat{x} \leq u \leq \hat{u}$. Therefore, \hat{x} is feasible for (B.2).

We have shown above that the feasibility of (B.1) implies the feasibility of (B.2). The inverse is also true by the same reasoning. Hence, we complete the proof. \square

Lemma B.3. *If two weighted bipartite graphs with vertex features corresponding to two LP problems are indistinguishable by the WL test, then these two problems share the same optimal objective value (could be ∞ or $-\infty$).*

Proof of Lemma B.3. If both problems are infeasible, then their optimal objective values are both ∞ . We then consider the case that both problems are feasible. We use the same setting and notations as in Lemma B.2, and in addition we have that $c_j = \hat{c}_j$, which is part of $h_j^W = \hat{h}_j^W$, is independent of $j \in J_q$ for any $q \in \{1, 2, \dots, t\}$. Suppose that x is an feasible solution to the problem (B.1) and let $\hat{x} \in \mathbb{R}^n$ be defined via $\hat{x}_j = \frac{1}{|J_q|} \sum_{j' \in J_q} x_{j'}$, $j \in J_q$. It is guaranteed by the proof of Lemma B.2 that \hat{x} is a feasible solution to (B.2). One can also see that $c^\top x = \hat{c}^\top \hat{x}$. Since this holds for any feasible solution x to (B.1), the optimal value of the objective function for (B.2) is smaller than or equal to that for (B.1). The inverse is also true and the proof is completed. \square

Lemma B.4. *Suppose that two weighted bipartite graphs with vertex features corresponding to two LP problems are indistinguishable by the WL test and their optimal objective values are both finite. Then these two problems have the same optimal solution with the smallest ℓ_2 -norm, up to permutation.*

Proof of Lemma B.4. We work with the same setting as in Lemma B.3, where permutations have already been applied. Let x and x' be the optimal solution to (B.1) and (B.2) with the smallest ℓ_2 -norm, respectively. (Recall that the optimal solution to a LP problem with the smallest ℓ_2 -norm is unique, see Remark 2.2.) Let $\hat{x} \in \mathbb{R}^n$ be defined via $\hat{x}_j = \frac{1}{|J_q|} \sum_{j' \in J_q} x_{j'}$ for $j \in J_q$, $q = 1, 2, \dots, t$. According to the arguments in the proof of Lemma B.2 and Lemma B.3, \hat{x} is an optimal solution to (B.2). The minimality of x' yields that

$$(B.3) \quad \|x'\|_2^2 \leq \|\hat{x}\|_2^2 = \sum_{q=1}^t |J_q| \left(\frac{1}{|J_q|} \sum_{j \in J_q} x_j \right)^2 = \sum_{q=1}^t \frac{1}{|J_q|} \left(\sum_{j \in J_q} x_j \right)^2 \leq \sum_{q=1}^t \sum_{j \in J_q} x_j^2 = \|x\|_2^2,$$

which implies $\|x'\| \leq \|x\|$. The converse $\|x\| \leq \|x'\|$ is also true. Therefore, we must have $\|x\| = \|x'\|$ and hence, the inequalities in (B.3) must hold as equalities. Then one can conclude that $x_j = x_{j'}$ for any $j, j' \in J_q$ and $q = 1, 2, \dots, t$, which leads to $x = \hat{x}$. Furthermore, it follows from $\|x'\| = \|\hat{x}\|$ and the uniqueness of x' (see Remark 2.2) that $x' = \hat{x} = x$, which completes the proof. \square

One corollary one can see from the proof of Lemma B.4 is that the components of the optimal solution with the smallest ℓ_2 -norm must be the same if the two corresponding vertices have the same color in the WL test.

Corollary B.5. *Let (G, H) be a weighted bipartite graph with vertex features and let x be the optimal solution to the corresponding LP problem with the smallest ℓ_2 -norm. Suppose that for some $j, j' \in \{1, 2, \dots, n\}$, one has $C_j^{l,W} = C_{j'}^{l,W}$ for any $l \in \mathbb{N}$ and any choices of hash functions, then $x_j = x_{j'}$.*

Let us also define another equivalence relationship on $\mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$ where colors of w_1, w_2, \dots, w_j with ordering (not just multisets) are considered:

Definition B.6. *Given $(G, H), (\hat{G}, \hat{H}) \in \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$, (G, H) and (\hat{G}, \hat{H}) are not in the same equivalence class of $\overset{W}{\sim}$ if and only if there exist some $L \in \mathbb{N}$ and some hash functions $HASH_{0,V}, HASH_{0,W}, HASH_{l,V}, HASH_{l,W}, HASH'_{l,V}$, and $HASH'_{l,W}$, $l = 1, 2, \dots, L$, such that $\{\{C_1^{L,V}, C_2^{L,V}, \dots, C_m^{L,V}\}\} \neq \{\{\hat{C}_1^{L,V}, \hat{C}_2^{L,V}, \dots, \hat{C}_m^{L,V}\}\}$ or $C_j^{l,W} \neq \hat{C}_j^{l,W}$ for some $j \in \{1, 2, \dots, n\}$.*

It is clear that $(G, H) \overset{W}{\sim} (\hat{G}, \hat{H})$ implies $(G, H) \sim (\hat{G}, \hat{H})$. One can actually obtain a stronger version of Lemma B.4 given $(G, H) \overset{W}{\sim} (\hat{G}, \hat{H})$.

Corollary B.7. *Suppose that two weighted bipartite graphs with vertex features corresponding to two LP problems, (G, H) and (\hat{G}, \hat{H}) , satisfies that $(G, H) \overset{W}{\sim} (\hat{G}, \hat{H})$. Then these two problems have the same optimal solution with the smallest ℓ_2 -norm.*

Proof of Corollary B.7. The proof of Lemma B.4 still applies with the difference that there is no permutation on $\{w_1, w_2, \dots, w_n\}$. \square

Proof of Theorem 4.1. The proof of Theorem 4.1 follows immediately from Lemmas B.2, B.3 and B.4. \square

APPENDIX C. SEPARATION POWER OF GRAPH NEURAL NETWORKS

This section aims to prove Theorem 4.2, i.e., the separation power of GNNs is equivalent to that of the WL test. Similar results can be found in previous literature, see e.g. [4, 19, 52]. We first introduce some lemmas that can directly imply Theorem 4.2. The lemma below, similar to [52, Lemma 2], states that the separation power of GNNs is at most that of the WL test.

Lemma C.1. *Let $(G, H), (\hat{G}, \hat{H}) \in \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$. If $(G, H) \sim (\hat{G}, \hat{H})$, then for any $F_W \in \mathcal{F}_{GNN}^W$, there exists a permutation $\sigma_W \in S_n$ such that $F_W(G, H) = \sigma_W(F_W(\hat{G}, \hat{H}))$.*

Proof of Lemma C.1. First we describe the sketch of our proof. The assumption $(G, H) \sim (\hat{G}, \hat{H})$ implies that, if we apply the WL test on (G, H) and (\hat{G}, \hat{H}) , the test results should be exactly the same whatever the hash functions in the WL test we choose. In the first step, we define a set of hash functions that are injective on all possible inputs. Second, we show that, if we apply an arbitrarily chosen GNN: $F_W \in \mathcal{F}_{GNN}^W$ on (G, H) and (\hat{G}, \hat{H}) , the vertex features of the two graphs are exactly the same up to permutation, given the fact that the WL test results

are the same. Finally, it concludes that $F_W(G, H)$ should be the same with $F_W(\hat{G}, \hat{H})$ up to permutation.

Let us first define hash functions. We choose $\text{HASH}_{0,V}$ and $\text{HASH}_{0,W}$ that are injective on the following sets (not multisets) respectively:

$$\{h_1^V, \dots, h_m^V, \hat{h}_1^V, \dots, \hat{h}_m^V\} \text{ and } \{h_1^W, \dots, h_n^W, \hat{h}_1^W, \dots, \hat{h}_n^W\}.$$

Let $\{C_i^{l-1,V}\}_{i=1}^m, \{C_j^{l-1,W}\}_{j=1}^n$ and $\{\hat{C}_i^{l-1,V}\}_{i=1}^m, \{\hat{C}_j^{l-1,W}\}_{j=1}^n$ be the vertex colors in the $(l-1)$ -th iteration ($1 \leq l \leq L$) in the WL test for (G, H) and (\hat{G}, \hat{H}) respectively. Define two sets (not multisets) that collect different colors:

$$\mathbf{C}_{l-1}^V = \{C_1^{l-1,V}, \dots, C_m^{l-1,V}, \hat{C}_1^{l-1,V}, \dots, \hat{C}_m^{l-1,V}\},$$

and

$$\mathbf{C}_{l-1}^W = \{C_1^{l-1,W}, \dots, C_n^{l-1,W}, \hat{C}_1^{l-1,W}, \dots, \hat{C}_n^{l-1,W}\}.$$

The hash function $\text{HASH}'_{l,V}$ and $\text{HASH}'_{l,W}$ are chosen such that the outputs are located in some linear spaces and that $\{\text{HASH}'_{l,V}(C) : C \in \mathbf{C}_{l-1}^V\}$ and $\{\text{HASH}'_{l,W}(C) : C \in \mathbf{C}_{l-1}^W\}$ are both linearly independent. Finally, we choose hash functions $\text{HASH}_{l,V}$ and $\text{HASH}_{l,W}$ such that $\text{HASH}_{l,V}$ is injective on the set (not multiset)

$$\left\{ \left(C_i^{l-1,V}, \sum_{j=1}^n E_{i,j} \text{HASH}'_{l,W} \left(C_j^{l-1,W} \right) \right) : 1 \leq i \leq m \right\} \\ \cup \left\{ \left(\hat{C}_i^{l-1,V}, \sum_{j=1}^n E_{i,j} \text{HASH}'_{l,W} \left(\hat{C}_j^{l-1,W} \right) \right) : 1 \leq i \leq m \right\},$$

and that $\text{HASH}_{l,W}$ is injective on the set (not multiset)

$$\left\{ \left(C_j^{l-1,W}, \sum_{i=1}^m E_{i,j} \text{HASH}'_{l,V} \left(C_i^{l-1,V} \right) \right) : 1 \leq j \leq n \right\} \\ \cup \left\{ \left(\hat{C}_j^{l-1,W}, \sum_{i=1}^m E_{i,j} \text{HASH}'_{l,V} \left(\hat{C}_i^{l-1,V} \right) \right) : 1 \leq j \leq n \right\}.$$

Those hash functions give the vertex colors at the next iteration (l -th layer): $\{C_i^{l,V}\}_{i=1}^m, \{C_j^{l,W}\}_{j=1}^n$ and $\{\hat{C}_i^{l,V}\}_{i=1}^m, \{\hat{C}_j^{l,W}\}_{j=1}^n$.

Consider any $F_W \in \mathcal{F}_{\text{GNN}}^W$ and let $\{h_i^{l-1,V}\}_{i=1}^m, \{h_j^{l-1,W}\}_{j=1}^n$ and $\{\hat{h}_i^{l-1,V}\}_{i=1}^m, \{\hat{h}_j^{l-1,W}\}_{j=1}^n$ be the vertex features in the l -th layer ($0 \leq l \leq L$) of the graph neural network F_W . (Update rule refers to equations (2.1),(2.2),(2.3),(2.5)) We aim to prove by induction that for any $l \in \{0, 1, \dots, L\}$, the followings hold:

- (i) $C_i^{l,V} = C_{i'}^{l,V}$ implies $h_i^{l,V} = h_{i'}^{l,V}$, for $1 \leq i, i' \leq m$;
- (ii) $\hat{C}_i^{l,V} = \hat{C}_{i'}^{l,V}$ implies $\hat{h}_i^{l,V} = \hat{h}_{i'}^{l,V}$, for $1 \leq i, i' \leq m$;
- (iii) $C_i^{l,V} = \hat{C}_{i'}^{l,V}$ implies $h_i^{l,V} = \hat{h}_{i'}^{l,V}$, for $1 \leq i, i' \leq m$;
- (iv) $C_j^{l,W} = C_{j'}^{l,W}$ implies $h_j^{l,W} = h_{j'}^{l,W}$, for $1 \leq j, j' \leq n$;
- (v) $\hat{C}_j^{l,W} = \hat{C}_{j'}^{l,W}$ implies $\hat{h}_j^{l,W} = \hat{h}_{j'}^{l,W}$, for $1 \leq j, j' \leq n$;

(vi) $C_j^{l,W} = \hat{C}_{j'}^{l,W}$ implies $h_j^{l,W} = \hat{h}_{j'}^{l,W}$, for $1 \leq j, j' \leq n$.

The above claims (i)-(vi) are clearly true for $l = 0$ due to the injectivity of $\text{HASH}_{0,V}$ and $\text{HASH}_{0,W}$. Now we assume that (i)-(vi) are true for some $l - 1 \in \{0, 1, \dots, L - 1\}$. Suppose that $C_i^{l,V} = C_{i'}^{l,V}$, i.e.,

$$\begin{aligned} \text{HASH}_{l,V} \left(C_i^{l-1,V}, \sum_{j=1}^n E_{i,j} \text{HASH}'_{l,W} \left(C_j^{l-1,W} \right) \right) \\ = \text{HASH}_{l,V} \left(C_{i'}^{l-1,V}, \sum_{j=1}^n E_{i',j} \text{HASH}'_{l,W} \left(C_j^{l-1,W} \right) \right), \end{aligned}$$

for some $1 \leq i, i' \leq m$. It follows from the injectivity of $\text{HASH}_{l,V}$ that

$$(C.1) \quad C_i^{l-1,V} = C_{i'}^{l-1,V},$$

and

$$\sum_{j=1}^n E_{i,j} \text{HASH}'_{l,W} \left(C_j^{l-1,W} \right) = \sum_{j=1}^n E_{i',j} \text{HASH}'_{l,W} \left(C_j^{l-1,W} \right).$$

According to the linearly independent property of $\text{HASH}'_{l,W}$, the above equation implies that

$$(C.2) \quad \sum_{C_j^{l-1,W}=C} E_{i,j} = \sum_{C_j^{l-1,W}=C} E_{i',j}, \quad \forall C \in \mathbf{C}_{l-1}^W.$$

Note that the induction assumption guarantees that $h_j^{l-1,W} = h_{j'}^{l-1,W}$ as long as $C_j^{l-1,W} = C_{j'}^{l-1,W}$. So one can assign for each $C \in \mathbf{C}_{l-1}^W$ some $h(C) \in \mathbb{R}^{d_{l-1}}$ such that $h_j^{l-1,W} = h(C)$ as long as $C_j^{l-1,W} = C$ for any $1 \leq j \leq n$. Therefore, it follows from (C.2) that

$$\begin{aligned} \sum_{j=1}^n E_{i,j} f_l^W(h_j^{l-1,W}) &= \sum_{C \in \mathbf{C}_{l-1}^W} \sum_{C_j^{l-1,W}=C} E_{i,j} f_l^W(h(C)) \\ &= \sum_{C \in \mathbf{C}_{l-1}^W} \sum_{C_j^{l-1,W}=C} E_{i',j} f_l^W(h(C)) = \sum_{j=1}^n E_{i',j} f_l^W(h_j^{l-1,W}). \end{aligned}$$

Note also that (C.1) and the induction assumption lead to $h_i^{l-1,V} = h_{i'}^{l-1,V}$. Then one can conclude that

$$h_i^{l,V} = g_l^V \left(h_i^{l-1,V}, \sum_{j=1}^n E_{i,j} f_l^W(h_j^{l-1,W}) \right) = g_l^V \left(h_{i'}^{l-1,V}, \sum_{j=1}^n E_{i',j} f_l^W(h_j^{l-1,W}) \right) = h_{i'}^{l,V}.$$

This proves the claim (i) for l . The other five claims can be proved using similar arguments.

Therefore, we obtain from $(G, H) \sim (\hat{G}, \hat{H})$ that

$$\left\{ \left\{ h_1^{L,V}, h_2^{L,V}, \dots, h_m^{L,V} \right\} \right\} = \left\{ \left\{ \hat{h}_1^{L,V}, \hat{h}_2^{L,V}, \dots, \hat{h}_m^{L,V} \right\} \right\},$$

and that

$$\left\{ \left\{ h_1^{L,W}, h_2^{L,W}, \dots, h_n^{L,W} \right\} \right\} = \left\{ \left\{ \hat{h}_1^{L,W}, \hat{h}_2^{L,W}, \dots, \hat{h}_n^{L,W} \right\} \right\}.$$

By the definition of the output layer, the above conclusion guarantees that $F_W(G, H) = \sigma_W(F_W(\hat{G}, \hat{H}))$ for some $\sigma_W \in S_n$. \square

Lemma C.2. *Let $(G, H), (\hat{G}, \hat{H}) \in \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$. Suppose that for any $F_W \in \mathcal{F}_{GNN}^W$, there exists a permutation $\sigma_W \in S_n$ such that $F_W(G, H) = \sigma_W(F_W(\hat{G}, \hat{H}))$. Then $F(G, H) = F(\hat{G}, \hat{H})$ holds for any $F \in \mathcal{F}_{GNN}$.*

Proof of Lemma C.2. Pick an arbitrary $F \in \mathcal{F}_{GNN}$. We choose $F_W \in \mathcal{F}_{GNN}^W$ such that

$$F_W(G', H') = (F(G', H'), \dots, F(G', H'))^\top \in \mathbb{R}^n, \quad \forall (G', H') \in \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W.$$

Note that every entry in the output of F_W is equal to the output of F . Thus, it follows from $F_W(G, H) = \sigma_W(F_W(\hat{G}, \hat{H}))$ that $F(G, H) = F(\hat{G}, \hat{H})$. \square

The next lemma is similar to [52, Theorem 3] and states that the separation power of GNNs is at least that of the WL test.

Lemma C.3. *Let $(G, H), (\hat{G}, \hat{H}) \in \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$. If $F(G, H) = F(\hat{G}, \hat{H})$ holds for any $F \in \mathcal{F}_{GNN}$, then $(G, H) \sim (\hat{G}, \hat{H})$.*

Proof of Lemma C.3. It suffices to prove that, if (G, H) can be distinguished from (\hat{G}, \hat{H}) by the WL test, then there exists $F \in \mathcal{F}_{GNN}$, such that $F(G, H) \neq F(\hat{G}, \hat{H})$. The distinguishability of the WL test implies that there exists $L \in \mathbb{N}$ and hash functions, $\text{HASH}_{0,V}$, $\text{HASH}_{0,W}$, $\text{HASH}_{l,V}$, $\text{HASH}_{l,W}$, $\text{HASH}'_{l,V}$, and $\text{HASH}'_{l,W}$, for $l = 1, 2, \dots, L$, such that

$$(C.3) \quad \left\{ \left\{ C_1^{L,V}, C_2^{L,V}, \dots, C_m^{L,V} \right\} \right\} \neq \left\{ \left\{ \hat{C}_1^{L,V}, \hat{C}_2^{L,V}, \dots, \hat{C}_m^{L,V} \right\} \right\},$$

or

$$(C.4) \quad \left\{ \left\{ C_1^{L,W}, C_2^{L,W}, \dots, C_n^{L,W} \right\} \right\} \neq \left\{ \left\{ \hat{C}_1^{L,W}, \hat{C}_2^{L,W}, \dots, \hat{C}_n^{L,W} \right\} \right\},$$

We aim to construct some GNNs such that the followings hold for any $l = 0, 1, \dots, L$:

- (i) $h_i^{l,V} = h_{i'}^{l,V}$ implies $C_i^{l,V} = C_{i'}^{l,V}$, for $1 \leq i, i' \leq m$;
- (ii) $\hat{h}_i^{l,V} = \hat{h}_{i'}^{l,V}$ implies $\hat{C}_i^{l,V} = \hat{C}_{i'}^{l,V}$, for $1 \leq i, i' \leq m$;
- (iii) $h_i^{l,V} = \hat{h}_{i'}^{l,V}$ implies $C_i^{l,V} = \hat{C}_{i'}^{l,V}$, for $1 \leq i, i' \leq m$;
- (iv) $h_j^{l,W} = h_{j'}^{l,W}$ implies $C_j^{l,W} = C_{j'}^{l,W}$, for $1 \leq j, j' \leq n$;
- (v) $\hat{h}_j^{l,W} = \hat{h}_{j'}^{l,W}$ implies $\hat{C}_j^{l,W} = \hat{C}_{j'}^{l,W}$, for $1 \leq j, j' \leq n$;
- (vi) $h_j^{l,W} = \hat{h}_{j'}^{l,W}$ implies $C_j^{l,W} = \hat{C}_{j'}^{l,W}$, for $1 \leq j, j' \leq n$.

It is clear that the above conditions (i)-(vi) hold for $l = 0$ as long as we choose f_{in}^V and f_{in}^W that are injective on the following two sets (not multisets) respectively:

$$\{h_1^V, \dots, h_m^V, \hat{h}_1^V, \dots, \hat{h}_m^V\} \text{ and } \{h_1^W, \dots, h_n^W, \hat{h}_1^W, \dots, \hat{h}_n^W\}.$$

We then assume that (i)-(vi) hold for some $0 \leq l-1 < L$, and show that these conditions are also satisfied for l if we choose $f_l^V, f_l^W, g_l^V, g_l^W$ properly. Let us consider the set (not multiset):

$$\{\alpha_1, \alpha_2, \dots, \alpha_s\} \subset \mathbb{R}^{d_{l-1}}$$

that collects all different values in $h_1^{l-1,W}, h_2^{l-1,W}, \dots, h_n^{l-1,W}, \hat{h}_1^{l-1,W}, \hat{h}_2^{l-1,W}, \dots, \hat{h}_n^{l-1,W}$. Let $d_l \geq s$ and let $e_p^{d_l} = (0, \dots, 0, 1, 0, \dots, 0)$ be the vector in \mathbb{R}^{d_l} with the p -th entry being 1 and

all other entries being 0, for $1 \leq p \leq s$. Choose $f_l^W : \mathbb{R}^{d_l-1} \rightarrow \mathbb{R}^{d_l}$ as a continuous function satisfying $f_l^W(\alpha_p) = e_p^{d_l}$, $p = 1, 2, \dots, s$, and choose $g_l^V : \mathbb{R}^{d_l-1} \times \mathbb{R}^{d_l} \rightarrow \mathbb{R}^{d_l}$ that is continuous and is injective when restricted on the set (not multiset)

$$\left\{ \left(h_i^{l-1,V}, \sum_{j=1}^n E_{i,j} f_l^W(h_j^{l-1,W}) \right) : 1 \leq i \leq m \right\} \\ \cup \left\{ \left(\hat{h}_i^{l-1,V}, \sum_{j=1}^n \hat{E}_{i,j} f_l^W(\hat{h}_j^{l-1,W}) \right) : 1 \leq i \leq m \right\}.$$

Noticing that

$$\sum_{j=1}^n E_{i,j} f_l^W(h_j^{l-1,W}) = \sum_{p=1}^s \left(\sum_{h_j^{l-1,W}=\alpha_p} E_{i,j} \right) e_p^{d_l},$$

and that $\{e_1^{d_l}, e_2^{d_l}, \dots, e_s^{d_l}\}$ is linearly independent, one can conclude that $h_i^{l,V} = h_i^{l,V}$ if and only if $h_i^{l-1,V} = h_i^{l-1,V}$ and $\sum_{j=1}^n E_{i,j} f_l^W(h_j^{l-1,W}) = \sum_{j=1}^n E_{i',j} f_l^W(h_j^{l-1,W})$, where the second condition is equivalent to

$$\sum_{h_j^{l-1,W}=\alpha_p} E_{i,j} = \sum_{h_j^{l-1,W}=\alpha_p} E_{i',j}, \quad \forall p \in \{1, 2, \dots, s\}.$$

This, as well as the condition (iv) for $l-1$, implies that

$$\sum_{j=1}^n E_{i,j} \text{HASH}'_{l,W}(C_j^{l-1,W}) = \sum_{j=1}^n E_{i',j} \text{HASH}'_{l,W}(C_j^{l-1,W}),$$

and hence that $C_i^{l,V} = C_{i'}^{l,V}$ by using $h_i^{l-1,V} = h_i^{l-1,V}$ and condition (i) for $l-1$. Therefore, we know that (i) is satisfied for l , and one can show (ii) and (iii) for l using similar arguments by taking d_l large enough. In addition, f_l^V and g_l^W can also be chosen in a similar way such that (iv)-(vi) are satisfied for l .

Combining (C.3), (C.4), and condition (i)-(iv) for L , we obtain that

$$(C.5) \quad \left\{ \left\{ h_1^{L,V}, h_2^{L,V}, \dots, h_m^{L,V} \right\} \right\} \neq \left\{ \left\{ \hat{h}_1^{L,V}, \hat{h}_2^{L,V}, \dots, \hat{h}_m^{L,V} \right\} \right\},$$

or

$$\left\{ \left\{ h_1^{L,W}, h_2^{L,W}, \dots, h_n^{L,W} \right\} \right\} \neq \left\{ \left\{ \hat{h}_1^{L,W}, \hat{h}_2^{L,W}, \dots, \hat{h}_n^{L,W} \right\} \right\}.$$

Without loss of generality, we can assume that (C.5) holds.

Consider the set (not multiset)

$$\{\beta_1, \beta_2, \dots, \beta_t\} \subset \mathbb{R}^{d_L},$$

that collects all different values in $h_1^{L,V}, h_2^{L,V}, \dots, h_m^{L,V}, \hat{h}_1^{L,V}, \hat{h}_2^{L,V}, \dots, \hat{h}_m^{L,V}$. Let $k > 1$ be a positive integer that is greater than the maximal multiplicity of an element in the multisets $\{\{h_1^{L,V}, h_2^{L,V}, \dots, h_m^{L,V}\}\}$ and $\{\{\hat{h}_1^{L,V}, \hat{h}_2^{L,V}, \dots, \hat{h}_m^{L,V}\}\}$. There exists a continuous function

$\varphi : \mathbb{R}^{d_L} \rightarrow \mathbb{R}$ such that $\varphi(\beta_q) = k^q$ for $q = 1, 2, \dots, t$, and due to (C.5) and the fact that the way of writing an integer as k -ary expression is unique, it hence holds that

$$\sum_{i=1}^m \varphi(h_i^{L,V}) \neq \sum_{i=1}^m \varphi(\hat{h}_i^{L,V}).$$

Set the dimension of $(L+1)$ -th layer as 1: $d_{L+1} = 1$, and set $f_{L+1,V} = 0$, $f_{L+1,W} = 0$, $g_{L+1,V}(h, 0) = \varphi(h)$, and $g_{L+1,W} = 0$. Then we have $h_i^{L+1,V} = \varphi(h_i^{L,V})$, $\hat{h}_i^{L+1,V} = \varphi(\hat{h}_i^{L,V})$, and $h_j^{L+1,W} = \hat{h}_j^{L+1,W}$ for $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, n$. Define $f_{\text{out}} : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ via $f_{\text{out}}(h, h') = h$. Then it follows that

$$\begin{aligned} f_{\text{out}} \left(\sum_{i=1}^m h_i^{L+1,V}, \sum_{j=1}^n h_j^{L+1,W} \right) &= \sum_{i=1}^m \varphi(h_i^{L,V}) \\ &\neq \sum_{i=1}^m \varphi(\hat{h}_i^{L,V}) = f_{\text{out}} \left(\sum_{i=1}^m \hat{h}_i^{L+1,V}, \sum_{j=1}^n \hat{h}_j^{L+1,W} \right), \end{aligned}$$

which guarantees the existence of $F \in \mathcal{F}_{\text{GNN}}$ that has $L+1$ layers and satisfies $F(G, H) \neq F(\hat{G}, \hat{H})$. \square

Proof of Theorem 4.2. The equivalence of the three conditions follow directly from Lemma C.1, C.2, and C.3. \square

Corollary C.4. *For any two weighted bipartite graphs with vertex features $(G, H), (\hat{G}, \hat{H}) \in \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$, the followings are equivalent:*

- (i) $(G, H) \stackrel{W}{\sim} (\hat{G}, \hat{H})$.
- (ii) For any $F_W \in \mathcal{F}_{\text{GNN}}^W$, it holds that $F_W(G, H) = F_W(\hat{G}, \hat{H})$.

Proof of Corollary C.4. The proof follows similar lines as in the proof of Theorem 4.2 with the difference that there is no permutation on $\{w_1, w_2, \dots, w_n\}$. \square

In addition to the separation power of GNNs for two weighted bipartite graphs with vertex features, one can also obtain results on separating different vertices in one weighted bipartite graph with vertex features.

Corollary C.5. *For any weighted bipartite graph with vertex features (G, H) and any $j, j' \in \{1, 2, \dots, n\}$, the followings are equivalent:*

- (i) $C_j^{l,W} = C_{j'}^{l,W}$ holds for any $l \in \mathbb{N}$ and any choice of hash functions.
- (ii) $F_W(G, H)_j = F_W(G, H)_{j'}, \forall F_W \in \mathcal{F}_{\text{GNN}}^W$.

Proof of Corollary C.5. “(i) \implies (ii)” and “(ii) \implies (i)” can be proved using similar arguments in the proof of Lemma C.1 and Lemma C.3, respectively. \square

APPENDIX D. UNIVERSAL APPROXIMATION OF \mathcal{F}_{GNN}

This section provides the proof of Theorem 4.3. The main mathematical tool used in the proof is the Stone–Weierstrass theorem:

Theorem D.1 (Stone–Weierstrass theorem [45, Section 5.7]). *Let X be a compact Hausdorff space and let $\mathcal{F} \subset \mathcal{C}(X, \mathbb{R})$ be a subalgebra. If \mathcal{F} separates points on X , i.e., for any $x, x' \in X$ with $x \neq x'$, there exists $F \in \mathcal{F}$ such that $F(x) \neq F(x')$, and $1 \in \mathcal{F}$, then \mathcal{F} is dense in $\mathcal{C}(X, \mathbb{R})$ with the topology of uniform convergence.*

Proof of Theorem 4.3. Let $\pi : X \rightarrow X/\sim$ be the quotient map, where $\pi(X) = X/\sim$ is equipped with the quotient topology. For any $F \in \mathcal{F}_{\text{GNN}}$, since $F : X \rightarrow \mathbb{R}$ is continuous and by Theorem 4.2, $F(G, H) = F(\hat{G}, \hat{H})$, $\forall (G, H) \sim (\hat{G}, \hat{H})$, there exists a unique continuous $\tilde{F} : \pi(X) \rightarrow \mathbb{R}$ such that $F = \tilde{F} \circ \pi$. Set

$$\tilde{\mathcal{F}}_{\text{GNN}} = \left\{ \tilde{F} : F \in \mathcal{F}_{\text{GNN}} \right\} \subset \mathcal{C}(\pi(X), \mathbb{R}).$$

In addition, the assumption on Φ , i.e.,

$$\Phi(G, H) = \Phi(\hat{G}, \hat{H}), \quad \forall (G, H) \sim (\hat{G}, \hat{H}),$$

leads to the existence of a unique $\tilde{\Phi} \in \mathcal{C}(\pi(X), \mathbb{R})$ with $\Phi = \tilde{\Phi} \circ \pi$.

Since X is compact, then $\pi(X)$ is also compact due to the continuity of π . According to Lemma D.2 below, $\tilde{\mathcal{F}}_{\text{GNN}}$ is a subalgebra of $\mathcal{C}(\pi(X), \mathbb{R})$. By Theorem 4.2, $\tilde{\mathcal{F}}_{\text{GNN}}$ separates points on $\pi(X)$. This can further imply that $\pi(X)$ is Hausdorff. In fact, for any $x, x' \in \pi(X)$, there exists $\tilde{F} \in \tilde{\mathcal{F}}_{\text{GNN}}$ with $\tilde{F}(x) \neq \tilde{F}(x')$. Without loss of generality, we assume that $\tilde{F}(x) < \tilde{F}(x')$ and choose some $c \in \mathbb{R}$ with $\tilde{F}(x) < c < \tilde{F}(x')$. By continuity of \tilde{F} , we know that $\tilde{F}^{-1}((-\infty, c)) \cap \pi(X)$ and $\tilde{F}^{-1}((c, +\infty)) \cap \pi(X)$ are disjoint open subsets of $\pi(X)$ with $x \in \tilde{F}^{-1}((-\infty, c)) \cap \pi(X)$ and $x' \in \tilde{F}^{-1}((c, +\infty)) \cap \pi(X)$, which leads to the Hausdorff property of $\pi(X)$. Note also that $1 \in \tilde{\mathcal{F}}_{\text{GNN}}$. Using Theorem D.1, we can conclude the denseness of $\tilde{\mathcal{F}}_{\text{GNN}}$ in $\mathcal{C}(\pi(X), \mathbb{R})$. Therefore, for any $\epsilon > 0$, there exists $F \in \mathcal{F}_{\text{GNN}}$, such that

$$\sup_{(G, H) \in X} |\Phi(G, H) - F(G, H)| = \sup_{x \in \pi(X)} |\tilde{\Phi}(x) - \tilde{F}(x)| < \epsilon,$$

which completes the proof. \square

Lemma D.2. \mathcal{F}_{GNN} is a subalgebra of $\mathcal{C}(\mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W, \mathbb{R})$, and as a corollary, $\tilde{\mathcal{F}}_{\text{GNN}}$ is a subalgebra of $\mathcal{C}(\mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W / \sim, \mathbb{R})$.

Proof of Lemma D.2. It suffices to show that \mathcal{F}_{GNN} is closed under addition and multiplication. Consider any $F, \hat{F} \in \mathcal{F}_{\text{GNN}}$. Thanks to Lemma D.3, we can assume that both F and \hat{F} have L layers. Suppose that F is constructed by

$$\begin{aligned} f_{\text{in}}^V : \mathcal{H}^V &\rightarrow \mathbb{R}^{d_0}, & f_{\text{in}}^W : \mathcal{H}^W &\rightarrow \mathbb{R}^{d_0}, \\ f_l^V, f_l^W : \mathbb{R}^{d_{l-1}} &\rightarrow \mathbb{R}^{d_l}, & g_l^V, g_l^W : \mathbb{R}^{d_{l-1}} \times \mathbb{R}^{d_l} &\rightarrow \mathbb{R}^{d_l}, \quad 1 \leq l \leq L, \\ f_{\text{out}} : \mathbb{R}^{d_L} \times \mathbb{R}^{d_L} &\rightarrow \mathbb{R}, \end{aligned}$$

and that \hat{F} is constructed by

$$\begin{aligned} \hat{f}_{\text{in}}^V : \mathcal{H}^V &\rightarrow \mathbb{R}^{\hat{d}_0}, & \hat{f}_{\text{in}}^W : \mathcal{H}^W &\rightarrow \mathbb{R}^{\hat{d}_0}, \\ \hat{f}_l^V, \hat{f}_l^W : \mathbb{R}^{\hat{d}_{l-1}} &\rightarrow \mathbb{R}^{\hat{d}_l}, & \hat{g}_l^V, \hat{g}_l^W : \mathbb{R}^{\hat{d}_{l-1}} \times \mathbb{R}^{\hat{d}_l} &\rightarrow \mathbb{R}^{\hat{d}_l}, \quad 1 \leq l \leq L, \\ \hat{f}_{\text{out}} : \mathbb{R}^{\hat{d}_L} \times \mathbb{R}^{\hat{d}_L} &\rightarrow \mathbb{R}. \end{aligned}$$

One can then construct two new GNNs computing $F + \hat{F}$ and $F \cdot \hat{F}$ as follows:

The input layer. The update rule of the input layer is defined by

$$\begin{aligned} \mathbf{f}_{\text{in}}^V : \mathcal{H}^V &\rightarrow \mathbb{R}^{d_0} \times \mathbb{R}^{\hat{d}_0}, \\ h &\mapsto \left(f_{\text{in}}^V(h), \hat{f}_{\text{in}}^V(h) \right), \\ \mathbf{f}_{\text{in}}^W : \mathcal{H}^W &\rightarrow \mathbb{R}^{d_0} \times \mathbb{R}^{\hat{d}_0}, \\ h &\mapsto \left(f_{\text{in}}^W(h), \hat{f}_{\text{in}}^W(h) \right). \end{aligned}$$

Then the vertex features after the computation of the input layer $\mathbf{h}_i^{0,V}$ and $\mathbf{h}_j^{0,W}$ are given by:

$$\begin{aligned} \mathbf{h}_i^{0,V} &= \mathbf{f}_{\text{in}}^V(h_i^V) = \left(f_{\text{in}}^V(h_i^V), \hat{f}_{\text{in}}^V(h_i^V) \right) = (h_i^{0,V}, \hat{h}_i^{0,V}) \in \mathbb{R}^{d_0} \times \mathbb{R}^{\hat{d}_0}, \\ \mathbf{h}_j^{0,W} &= \mathbf{f}_{\text{in}}^W(h_j^W) = \left(f_{\text{in}}^W(h_j^W), \hat{f}_{\text{in}}^W(h_j^W) \right) = (h_j^{0,W}, \hat{h}_j^{0,W}) \in \mathbb{R}^{d_0} \times \mathbb{R}^{\hat{d}_0}, \end{aligned}$$

for $i = 1, 2, \dots, m$, and $j = 1, 2, \dots, n$.

The l -th layer ($1 \leq l \leq L$). We set

$$\begin{aligned} \mathbf{f}_l^V : \mathbb{R}^{d_{l-1}} \times \mathbb{R}^{\hat{d}_{l-1}} &\rightarrow \mathbb{R}^{d_l} \times \mathbb{R}^{\hat{d}_l}, \\ (h, \hat{h}) &\mapsto \left(f_l^V(h), \hat{f}_l^V(\hat{h}) \right), \\ \mathbf{f}_l^W : \mathbb{R}^{d_{l-1}} \times \mathbb{R}^{\hat{d}_{l-1}} &\rightarrow \mathbb{R}^{d_l} \times \mathbb{R}^{\hat{d}_l}, \\ (h, \hat{h}) &\mapsto \left(f_l^W(h), \hat{f}_l^W(\hat{h}) \right), \\ \mathbf{g}_l^V : \mathbb{R}^{d_{l-1}} \times \mathbb{R}^{\hat{d}_{l-1}} \times \mathbb{R}^{d_l} \times \mathbb{R}^{\hat{d}_l} &\rightarrow \mathbb{R}^{d_l} \times \mathbb{R}^{\hat{d}_l}, \\ (h, \hat{h}, h', \hat{h}') &\mapsto \left(g_l^V(h, h'), \hat{g}_l^V(\hat{h}, \hat{h}') \right), \end{aligned}$$

and

$$\begin{aligned} \mathbf{g}_l^W : \mathbb{R}^{d_{l-1}} \times \mathbb{R}^{\hat{d}_{l-1}} \times \mathbb{R}^{d_l} \times \mathbb{R}^{\hat{d}_l} &\rightarrow \mathbb{R}^{d_l} \times \mathbb{R}^{\hat{d}_l}, \\ (h, \hat{h}, h', \hat{h}') &\mapsto \left(g_l^W(h, h'), \hat{g}_l^W(\hat{h}, \hat{h}') \right). \end{aligned}$$

Then the vertex features after the computation of the l -th layer $\mathbf{h}_i^{l,V}$ and $\mathbf{h}_j^{l,W}$ are given by:

$$\begin{aligned} \mathbf{h}_i^{l,V} &= \mathbf{g}_l^V \left(\mathbf{h}_i^{l-1,V}, \sum_{j=1}^n E_{i,j} \mathbf{f}_l^W(\mathbf{h}_j^{l-1,W}) \right) \\ &= \left(g_l^V \left(h_i^{l-1,V}, \sum_{j=1}^n E_{i,j} f_l^W(h_j^{l-1,W}) \right), \hat{g}_l^V \left(\hat{h}_i^{l-1,V}, \sum_{j=1}^n E_{i,j} \hat{f}_l^W(\hat{h}_j^{l-1,W}) \right) \right) \\ &= (h_i^{l,V}, \hat{h}_i^{l,V}) \in \mathbb{R}^{d_l} \times \mathbb{R}^{\hat{d}_l}, \end{aligned}$$

and

$$\mathbf{h}_j^{l,W} = \mathbf{g}_l^W \left(\mathbf{h}_j^{l-1,W}, \sum_{i=1}^m E_{i,j} \mathbf{f}_l^V(\mathbf{h}_i^{l-1,V}) \right)$$

$$\begin{aligned}
&= \left(g_l^W \left(h_j^{l-1, W}, \sum_{i=1}^m E_{i,j} f_l^V(h_i^{l-1, V}) \right), \hat{g}_l^W \left(\hat{h}_j^{l-1, W}, \sum_{i=1}^m E_{i,j} \hat{f}_l^V(\hat{h}_i^{l-1, V}) \right) \right) \\
&= (h_j^{l, W}, \hat{h}_j^{l, W}) \in \mathbb{R}^{d_l} \times \mathbb{R}^{\hat{d}_l},
\end{aligned}$$

for $i = 1, 2, \dots, m$, and $j = 1, 2, \dots, n$.

The output layer. To obtain $F + \hat{F}$, we set

$$\begin{aligned}
\mathbf{f}_{\text{out}}^{\text{add}} : \mathbb{R}^{d_L} \times \mathbb{R}^{\hat{d}_L} \times \mathbb{R}^{d_L} \times \mathbb{R}^{\hat{d}_L} &\rightarrow \mathbb{R}, \\
((h, \hat{h}), (h', \hat{h}')) &\mapsto f_{\text{out}}(h, h') + \hat{f}_{\text{out}}(\hat{h}, \hat{h}').
\end{aligned}$$

Then it holds that

$$\begin{aligned}
\mathbf{f}_{\text{out}}^{\text{add}} \left(\sum_{i=1}^m \mathbf{h}_i^{L, V}, \sum_{j=1}^n \mathbf{h}_j^{L, W} \right) &= f_{\text{out}} \left(\sum_{i=1}^m h_i^{L, V}, \sum_{j=1}^n h_j^{L, W} \right) + \hat{f}_{\text{out}} \left(\sum_{i=1}^m \hat{h}_i^{L, V}, \sum_{j=1}^n \hat{h}_j^{L, W} \right) \\
&= F(G, H) + \hat{F}(G, H).
\end{aligned}$$

To obtain $F \cdot \hat{F}$, we set

$$\begin{aligned}
\mathbf{f}_{\text{out}}^{\text{multiply}} : \mathbb{R}^{d_L} \times \mathbb{R}^{\hat{d}_L} \times \mathbb{R}^{d_L} \times \mathbb{R}^{\hat{d}_L} &\rightarrow \mathbb{R}, \\
((h, \hat{h}), (h', \hat{h}')) &\mapsto f_{\text{out}}(h, h') \cdot \hat{f}_{\text{out}}(\hat{h}, \hat{h}').
\end{aligned}$$

Then it holds that

$$\begin{aligned}
\mathbf{f}_{\text{out}}^{\text{multiply}} \left(\sum_{i=1}^m \mathbf{h}_i^{L, V}, \sum_{j=1}^n \mathbf{h}_j^{L, W} \right) &= f_{\text{out}} \left(\sum_{i=1}^m h_i^{L, V}, \sum_{j=1}^n h_j^{L, W} \right) \cdot \hat{f}_{\text{out}} \left(\sum_{i=1}^m \hat{h}_i^{L, V}, \sum_{j=1}^n \hat{h}_j^{L, W} \right) \\
&= F(G, H) \cdot \hat{F}(G, H).
\end{aligned}$$

The constructed GNNs satisfy $F(G, H) + \hat{F}(G, H) \in \mathcal{F}_{\text{GNN}}$ and $F(G, H) \cdot \hat{F}(G, H) \in \mathcal{F}_{\text{GNN}}$, which finishes the proof. \square

Lemma D.3. *If $F \in \mathcal{F}_{\text{GNN}}$ has L layers, then there exists $\hat{F} \in \mathcal{F}_{\text{GNN}}$ with $L + 1$ layers such that $F = \hat{F}$.*

Proof of Lemma D.3. Suppose that F is constructed by $f_{\text{in}}^V, f_{\text{in}}^W, f_{\text{out}}, \{f_l^V, f_l^W, g_l^V, g_l^W\}_{l=0}^L$. We choose $f_{L+1}^V \equiv 0, f_{L+1}^W \equiv 0, g_{L+1}^V(h, h') = h, g_{L+1}^W(h, h') = h$. Let \hat{F} be constructed by $f_{\text{in}}^V, f_{\text{in}}^W, f_{\text{out}}, \{f_l^V, f_l^W, g_l^V, g_l^W\}_{l=0}^{L+1}$. Then \hat{F} has $L + 1$ layers with $\hat{F} = F$. \square

APPENDIX E. UNIVERSAL APPROXIMATION OF $\mathcal{F}_{\text{GNN}}^W$

This section provides an universal approximation result of $\mathcal{F}_{\text{GNN}}^W$.

Theorem E.1. *Let $X \subset \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$ be a compact subset that is closed under the action of $S_m \times S_n$. Suppose that $\Phi \in \mathcal{C}(X, \mathbb{R}^n)$ satisfies the followings:*

(i) *For any $\sigma_V \in S_m, \sigma_W \in S_n$, and $(G, H) \in X$, it holds that*

$$(E.1) \quad \Phi((\sigma_V, \sigma_W) * (G, H)) = \sigma_W(\Phi(G, H)).$$

(ii) $\Phi(G, H) = \Phi(\hat{G}, \hat{H})$ holds for all $(G, H), (\hat{G}, \hat{H}) \in X$ with $(G, H) \stackrel{W}{\sim} (\hat{G}, \hat{H})$.

(iii) Given any $(G, H) \in X$ and any $j, j' \in \{1, 2, \dots, n\}$, if $C_j^{l,W} = C_{j'}^{l,W}$ (the vertex colors obtained in the l -th iteration in WL test) holds for any $l \in \mathbb{N}$ and any choices of hash functions, then $\Phi(G, H)_j = \Phi(G, H)_{j'}$.

Then for any $\epsilon > 0$, there exists $F \in \mathcal{F}_{\text{GNN}}^W$ such that

$$\sup_{(G,H) \in X} \|\Phi(G, H) - F(G, H)\| < \epsilon.$$

Theorem E.1 is a LP-graph version of results on the closure of equivariant GNN class in [4, 19]. The main tool in the proof of Theorem E.1 is the following generalized Stone-Weierstrass theorem for equivariant functions established in [4].

Theorem E.2 (Generalized Stone-Weierstrass theorem [4, Theorem 22]). *Let X be a compact topology space and let \mathbf{G} be a finite group that acts continuously on X and \mathbb{R}^n . Define the collection of all equivariant continuous functions from X to \mathbb{R}^n as follows:*

$$\mathcal{C}_E(X, \mathbb{R}^n) = \{F \in \mathcal{C}(X, \mathbb{R}^n) : F(g * x) = g * F(x), \forall x \in X, g \in \mathbf{G}\}.$$

Consider any $\mathcal{F} \subset \mathcal{C}_E(X, \mathbb{R}^n)$ and any $\Phi \in \mathcal{C}_E(X, \mathbb{R}^n)$. Suppose the following conditions hold:

- (i) \mathcal{F} is a subalgebra of $\mathcal{C}(X, \mathbb{R}^n)$ and $\mathbf{1} \in \mathcal{F}$.
- (ii) For any $x, x' \in X$, if $f(x) = f(x')$ holds for any $f \in \mathcal{C}(X, \mathbb{R})$ with $f\mathbf{1} \in \mathcal{F}$, then for any $F \in \mathcal{F}$, there exists $g \in \mathbf{G}$ such that $F(x) = g * F(x')$.
- (iii) For any $x, x' \in X$, if $F(x) = F(x')$ holds for any $F \in \mathcal{F}$, then $\Phi(x) = \Phi(x')$.
- (iv) For any $x \in X$, it holds that $\Phi(x)_j = \Phi(x)_{j'}, \forall (j, j') \in J(x)$, where

$$J(x) = \{\{1, 2, \dots, n\}^n : F(x)_j = F(x)_{j'}, \forall F \in \mathcal{F}\}.$$

Then for any $\epsilon > 0$, there exists $F \in \mathcal{F}$ such that

$$\sup_{x \in X} \|\Phi(x) - F(x)\| < \epsilon.$$

We refer to [49] for different versions of Stone-Weierstrass theorem, that is also used in [4]. In the proof of Theorem E.1, we also need the following lemma whose proof is almost the same as the proof of Lemma D.2 and is hence omitted.

Lemma E.3. $\mathcal{F}_{\text{GNN}}^W$ is a subalgebra of $\mathcal{C}(\mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W, \mathbb{R}^n)$.

Proof of Theorem E.1. Let $S_m \times S_n$ act on \mathbb{R}^n via

$$(\sigma_V, \sigma_W) * y = \sigma_W(y), \quad \forall \sigma_V \in S_m, \sigma_W \in S_n, y \in \mathbb{R}^n.$$

Then it follows from (E.1) that Φ is equivariant. In addition, the definition of graph neural networks directly guarantees the equivariance of functions in $\mathcal{F}_{\text{GNN}}^W$. Therefore, one only needs to verify the conditions with \mathcal{F} as $\mathcal{F}_{\text{GNN}}^W$ and \mathbf{G} as S_n in Theorem E.2:

- Condition (i) in Theorem E.2 follows from Lemma E.3 and the definition of $\mathcal{F}_{\text{GNN}}^W$.
- Condition (ii) in Theorem E.2 follows from Theorem 4.2 and $\mathcal{F}_{\text{GNN}}^W \mathbf{1} \subset \mathcal{F}_{\text{GNN}}^W$.
- Condition (iii) in Theorem E.2 follows from Corollary C.4 and Condition (ii) in Theorem E.1.
- Condition (iv) in Theorem E.2 follows from Corollary C.5 and Condition (iii) in Theorem E.1.

It finishes the proof. \square

APPENDIX F. PROOF OF MAIN THEOREMS

We collect the proofs of main theorems stated in Section 3 in this section.

Proof of Theorem 3.1. If $F(G, H) = F(\hat{G}, \hat{H})$ holds for any $F \in \mathcal{F}_{\text{GNN}}$, then Theorem 4.2 guarantees that $(G, H) \sim (\hat{G}, \hat{H})$. Thus, Condition (i), (ii), and (iii) follow directly from Lemmas B.2, B.3, and B.4, respectively.

Furthermore, if $F_W(G, H) = F_W(\hat{G}, \hat{H})$, $\forall F_W \in \mathcal{F}_{\text{GNN}}^W$, then it follows from Corollary C.4 and Corollary B.7 that the two LP problems associated to (G, H) and (\hat{G}, \hat{H}) share the same optimal solution with the smallest ℓ_2 -norm. \square

Then we head into the proof of three main approximation theorems, say Theorem 3.2, 3.4, and 3.6, that state that GNNs can approximate the feasibility mapping Φ_{feas} , the optimal objective value mapping Φ_{obj} , and the optimal solution mapping Φ_{solu} with arbitrarily small error, respectively. We have established in Sections D and E several theorems for graph neural networks to approximate continuous mappings. Therefore, the proof of Theorem 3.2, 3.4, and 3.6 basically consists of two steps:

- (i) Show that the mappings Φ_{feas} , Φ_{obj} , and Φ_{solu} are measurable.
- (ii) Use continuous mappings to approximate the target measurable mappings, and then apply the universal approximation results established in Sections D and E.

Let us first prove the feasibility of Φ_{feas} , Φ_{obj} , and Φ_{solu} in the following three lemmas.

Lemma F.1. *The feasibility mapping Φ_{feas} defined in (2.7) is measurable, i.e., the preimages $\Phi_{\text{feas}}^{-1}(0)$ and $\Phi_{\text{feas}}^{-1}(1)$ are both measurable subsets of $\mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$.*

Proof of Lemma F.1. It suffices to prove that for any $\circ \in \{\leq, =, \geq\}^m$ and any $N_l, N_u \subset \{1, 2, \dots, n\}$, the set

$$X_{\text{feas}} := \{(A, b, l, u) \in \mathbb{R}^{m \times n} \times \mathbb{R}^m \times \mathbb{R}^{|N_l|} \times \mathbb{R}^{|N_u|} : \\ \exists x \in \mathbb{R}^n, \text{ s.t. } Ax \circ b, x_j \geq l_j, \forall j \in N_l, x_j \leq u_j, \forall j \in N_u\},$$

is a measurable subset in $\mathbb{R}^{m \times n} \times \mathbb{R}^m \times \mathbb{R}^{|N_l|} \times \mathbb{R}^{|N_u|}$. Without loss of generality, we assume that $\circ = (\leq, \dots, \leq, =, \dots, =, \geq, \dots, \geq)$ where “ \leq ”, “ $=$ ”, and “ \geq ” appear for k_1 , $k_2 - k_1$, and $m - k_1 - k_2$ times, respectively, $0 \leq k_1 \leq k_2 \leq m$.

Let us define a function $V_{\text{feas}} : \mathbb{R}^{m \times n} \times \mathbb{R}^m \times \mathbb{R}^{|N_l|} \times \mathbb{R}^{|N_u|} \times \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$ that measures to what extend a point in \mathbb{R}^n violates the constraints:

$$V_{\text{feas}}(A, b, l, u, x) = \max \left\{ \max_{1 \leq i \leq k_1} \left(\sum_{j=1}^n A_{i,j} x_j - b_i \right)_+, \max_{k_1 < i \leq k_2} \left| \sum_{j=1}^n A_{i,j} x_j - b_i \right|, \right. \\ \left. \max_{k_2 < i \leq m} \left(b_i - \sum_{j=1}^n A_{i,j} x_j \right)_+, \max_{j \in N_l} (l_j - x_j)_+, \max_{j \in N_u} (x_j - u_j)_+ \right\},$$

where $y_+ = \max\{y, 0\}$. It can be seen that V_{feas} is continuous and $V_{\text{feas}}(A, b, l, u, x) = 0$ if and only if $Ax \circ b$, $x_j \geq l_j$, $\forall j \in N_l$, and $x_j \leq u_j$, $\forall j \in N_u$. Therefore, for any $(A, b, l, u) \in \mathbb{R}^{m \times n} \times \mathbb{R}^m \times \mathbb{R}^{|N_l|} \times \mathbb{R}^{|N_u|}$, the followings are equivalent:

- $(A, b, l, u) \in X_{\text{feas}}$.
- There exists $R \in \mathbb{N}_+$ and $x \in B_R := \{x' \in \mathbb{R}^n : \|x'\| \leq R\}$, such that $V_{\text{feas}}(A, b, l, u, x) = 0$.
- There exists $R \in \mathbb{N}_+$ such that for any $r \in \mathbb{N}_+$, $V_{\text{feas}}(A, b, l, u, x) \leq 1/r$ holds for some $x \in B_R \cap \mathbb{Q}^n$.

This implies that X_{feas} can be described via

$$\bigcup_{R \in \mathbb{N}_+} \bigcap_{r \in \mathbb{N}_+} \bigcup_{x \in B_R \cap \mathbb{Q}^n} \left\{ (A, b, l, u) \in \mathbb{R}^{m \times n} \times \mathbb{R}^m \times \mathbb{R}^{|N_l|} \times \mathbb{R}^{|N_u|} : V_{\text{feas}}(A, b, l, u, x) \leq \frac{1}{r} \right\}.$$

Since \mathbb{Q}^n is countable and V is continuous, we immediately obtain from the above expression that X_{feas} is measurable. \square

Lemma F.2. *The optimal objective value mapping Φ_{obj} defined in (2.8) is measurable.*

Proof of Lemma F.2. It suffices to prove that for any $\circ \in \{\leq, \geq\}^m$, any subsets $N_l, N_u \subset \{1, 2, \dots, n\}$, and any $\phi \in \mathbb{R}$, the set

$$X_{\text{obj}} := \{(A, b, c, l, u) \in \mathbb{R}^{m \times n} \times \mathbb{R}^m \times \mathbb{R}^n \times \mathbb{R}^{|N_l|} \times \mathbb{R}^{|N_u|} : \\ \exists x \in \mathbb{R}^n, \text{ s.t. } c^\top x \leq \phi, Ax \circ b, x_j \geq l_j, \forall j \in N_l, x_j \leq u_j, \forall j \in N_u\},$$

is a measurable subset in $\mathbb{R}^{m \times n} \times \mathbb{R}^m \times \mathbb{R}^n \times \mathbb{R}^{|N_l|} \times \mathbb{R}^{|N_u|}$. Then the proof follows the same lines as in the proof of Lemma F.1, with a different violation function

$$V_{\text{obj}}(A, b, c, l, u, x) = \max\{(c^\top x - \phi)_+, V_{\text{feas}}(A, b, l, u, x)\}.$$

\square

Lemma F.3. *The optimal solution mapping Φ_{solu} defined in (2.9) is measurable.*

Proof of Lemma F.3. It suffices to show that for every $j_0 \in \{1, 2, \dots, n\}$, the mapping

$$\pi_{j_0} \circ \Phi_{\text{solu}} : \Phi_{\text{obj}}^{-1}(\mathbb{R}) \rightarrow \mathbb{R},$$

is measurable, where $\pi_{j_0} : \mathbb{R}^n \rightarrow \mathbb{R}$ maps a vector $x \in \mathbb{R}^n$ to its j_0 -th component. Similar as before, one can consider any $\circ \in \{\leq, \geq\}^m$, any $N_l, N_u \subset \{1, 2, \dots, n\}$, and any $\phi \in \mathbb{R}$, and prove that the set

$$X_{\text{solu}} := \{(A, b, c, l, u) \in \mathbb{R}^{m \times n} \times \mathbb{R}^m \times \mathbb{R}^n \times \mathbb{R}^{|N_l|} \times \mathbb{R}^{|N_u|} :$$

$$\text{The LP problem, } \min_{x \in \mathbb{R}^d} c^\top x, \text{ s.t. } Ax \circ b, x_j \geq l_j, \forall j \in N_l, x_j \leq u_j, \forall j \in N_u,$$

has a finite optimal objective value, and its optimal solution with the smallest ℓ_2 -norm,

$$x_{\text{opt}}, \text{ satisfies } (x_{\text{opt}})_{j_0} < \phi\},$$

is measurable.

Note that we have fixed $\circ \in \{\leq, \geq\}^m$ and $N_l, N_u \subset \{1, 2, \dots, n\}$. Let

$$\iota : \mathbb{R}^{m \times n} \times \mathbb{R}^m \times \mathbb{R}^n \times \mathbb{R}^{|N_l|} \times \mathbb{R}^{|N_u|} \rightarrow \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W,$$

be the embedding map. Define another violation function

$$V_{\text{solu}} : (\Phi_{\text{obj}} \circ \iota)^{-1}(\mathbb{R}) \times \mathbb{R}^n \rightarrow \mathbb{R},$$

via

$$V_{\text{solu}}(A, b, c, l, u, x) = \max \left\{ (c^\top x - \Phi_{\text{obj}}(\iota(A, b, c, l, u)))_+, V_{\text{feas}}(A, b, c, l, u, x) \right\},$$

which is measurable due to the measurability of Φ_{obj} and the continuity of V_{feas} . Moreover, V_{solu} is continuous with respect to x . Therefore, the followings are equivalent for $(A, b, c, l, u) \in (\Phi_{\text{obj}} \circ \iota)^{-1}(\mathbb{R})$:

- $(A, b, c, l, u) \in X_{\text{solu}}$.
- There exists $x \in \mathbb{R}^n$ with $x_{j_0} < \phi$, such that $V_{\text{solu}}(A, b, c, l, u, x) = 0$ and $V_{\text{solu}}(A, b, c, l, u, x') > 0$, $\forall x' \in B_{\|x\|}$, $x'_{j_0} \geq \phi$.
- There exists $R \in \mathbb{Q}_+$, $r \in \mathbb{N}_+$, and $x \in B_R$ with $x_{j_0} \leq \phi - 1/r$, such that $V_{\text{solu}}(A, b, c, l, u, x) = 0$ and $V_{\text{solu}}(A, b, c, l, u, x') > 0$, $\forall x' \in B_R$, $x'_{j_0} \geq \phi$.
- There exists $R \in \mathbb{Q}_+$ and $r \in \mathbb{N}_+$, such that for all $r' \in \mathbb{N}_+$, $\exists x \in B_R \cap \mathbb{Q}^n$, $x_{j_0} \leq \phi - 1/r$, s.t. $V_{\text{solu}}(A, b, c, l, u, x) < 1/r'$ and that $\exists r'' \in \mathbb{N}_+$, s.t., $V_{\text{solu}}(A, b, c, l, u, x') \geq 1/r''$, $\forall x' \in B_R \cap \mathbb{Q}^n$, $x'_{j_0} \geq \phi$.

Therefore, one has that

$$\begin{aligned} X_{\text{solu}} &= \bigcup_{R \in \mathbb{Q}_+} \bigcup_{r \in \mathbb{N}_+} \\ &\left(\left(\bigcap_{r' \in \mathbb{N}_+} \bigcup_{x \in B_R \cap \mathbb{Q}^n, x_{j_0} \leq \phi - \frac{1}{r}} \left\{ (A, b, c, l, u) \in (\Phi_{\text{obj}} \circ \iota)^{-1}(\mathbb{R}) : V_{\text{solu}}(A, b, c, l, u, x) < \frac{1}{r'} \right\} \right) \right. \\ &\left. \cap \left(\bigcup_{r'' \in \mathbb{N}_+} \bigcap_{x' \in B_R \cap \mathbb{Q}^n, x'_{j_0} \geq \phi} \left\{ (A, b, c, l, u) \in (\Phi_{\text{obj}} \circ \iota)^{-1}(\mathbb{R}) : V_{\text{solu}}(A, b, c, l, u, x') \geq \frac{1}{r''} \right\} \right) \right), \end{aligned}$$

which is measurable. \square

With the measurability of Φ_{feas} , Φ_{obj} , and Φ_{solu} established, the next step is to approximate Φ_{feas} , Φ_{obj} , and Φ_{solu} using continuous mappings, and hence graph neural networks. Before proceeding, let us mention that $\mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$ is essentially the disjoint union of finitely many product spaces of Euclidean spaces and discrete spaces that have finitely many points and are equipped with discrete measures. More specifically,

$$\begin{aligned} \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W &\cong \mathbb{R}^{m \times n} \times (\mathbb{R} \times \{\leq, =, \geq\})^m \times (\mathbb{R} \times (\mathbb{R} \cup \{-\infty\})) \times (\mathbb{R} \cup \{+\infty\})^n \\ &\cong \bigcup_{k,k'=0}^n \bigcup_{s=1}^{\binom{n}{k}} \bigcup_{s'=1}^{\binom{n}{k'}} \mathbb{R}^{m \times n} \times \mathbb{R}^m \times \mathbb{R}^n \times \mathbb{R}^{n-k} \times \mathbb{R}^{n-k'} \\ &\quad \times \{\leq, =, \geq\}^m \times \{-\infty\}^k \times \{+\infty\}^{k'}. \end{aligned}$$

Therefore, many results in real analysis for Euclidean spaces still apply for $\mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$ and $\text{Meas}(\cdot)$, including the following Lusin's theorem.

Theorem F.4 (Lusin's theorem [17, Theorem 1.14]). *Let μ be a Borel regular measure on \mathbb{R}^n and let $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be μ -measurable. Then for any μ -measurable $X \subset \mathbb{R}^n$ with $\mu(X) < \infty$ and any $\epsilon > 0$, there exists a compact set $E \subset X$ with $\mu(X \setminus E) < \epsilon$, such that $f|_E$ is continuous.*

Proof of Theorem 3.2. Since $X \subset \mathcal{G}_{m,n} \times \mathcal{H}_m^V \times \mathcal{H}_n^W$ is measurable with finite measure, according to Lusin's theorem, there is a compact set $E \subset X$ such that $\Phi_{\text{feas}}|_E$ is continuous with $\text{Meas}(X \setminus E) < \epsilon$. By Lemma B.2 and Theorem 4.3, there exists $F \in \mathcal{F}_{\text{GNN}}$ such that

$$\sup_{(G,H) \in E} |F(G, H) - \Phi_{\text{feas}}(G, H)| < \frac{1}{2},$$

which implies that

$$\mathbb{1}_{F(G,H) > 1/2} = \Phi_{\text{feas}}(G, H), \quad \forall (G, H) \in E.$$

Therefore, one obtains

$$\text{Meas}(\{(G, H) \in X : \mathbb{1}_{F(G,H) > 1/2} \neq \Phi_{\text{feas}}(G, H)\}) \leq \text{Meas}(X \setminus E) < \epsilon,$$

and the proof is completed. \square

Proof of Corollary 3.3. As a finite set, \mathcal{D} is compact and $\Phi_{\text{feas}}|_{\mathcal{D}}$ is continuous. The rest of the proof is similar to that of Theorem 3.2, using Lemma B.2 and Theorem 4.3. \square

Proof of Theorem 3.4. (i) The proof follows the same lines as in the proof of Theorem 3.2, with the difference that we approximate

$$\Phi(G, H) = \begin{cases} 1, & \text{if } \Phi_{\text{obj}}(G, H) \in \mathbb{R}, \\ 0, & \text{otherwise,} \end{cases}$$

instead of Φ_{feas} and we use Lemma B.3 instead of Lemma B.2.

(ii) The proof is still similar to that of Theorem 3.2. By Lusin's theorem, there is a compact set $E \subset X \cap \Phi_{\text{obj}}^{-1}(\mathbb{R})$ such that $\Phi_{\text{obj}}|_E$ is continuous with $\text{Meas}((X \cap \Phi_{\text{obj}}^{-1}(\mathbb{R})) \setminus E) < \epsilon$. Using Lemma B.3 and Theorem 4.3, there exists $F_2 \in \mathcal{F}_{\text{GNN}}$ with

$$\sup_{(G,H) \in E} |F_2(G, H) - \Phi_{\text{obj}}(G, H)| < \delta,$$

which implies that

$$\text{Meas}(\{(G, H) \in X : |F_2(G, H) - \Phi_{\text{obj}}(G, H)| > \delta\}) \leq \text{Meas}((X \cap \Phi_{\text{obj}}^{-1}(\mathbb{R})) \setminus E) < \epsilon.$$

\square

Proof of Corollary 3.5. The results can be proved using similar techniques as in Theorem 3.2 and Theorem 3.4 by noticing that any finite dataset is compact on which any real-valued function is continuous. \square

Proof of Theorem 3.6. By Lusin's theorem, there exists a compact subset $E' \subset X$ such that $\text{Meas}(A \setminus X) < \epsilon/|S_m \times S_n|$ and that $\Phi_{\text{solu}}|_{E'}$ is continuous. Define another compact set:

$$E = \bigcap_{(\sigma_V, \sigma_W) \in S_m \times S_n} (\sigma_V, \sigma_W) * E' \subset X,$$

which is closed under the action of $S_m \times S_n$ and satisfies

$$\begin{aligned}
\text{Meas}(X \setminus E) &\leq \sum_{(\sigma_V, \sigma_W) \in S_m \times S_n} \text{Meas}(X \setminus (\sigma_V, \sigma_W) * E') \\
&= \sum_{(\sigma_V, \sigma_W) \in S_m \times S_n} \text{Meas}((\sigma_V, \sigma_W) * X \setminus (\sigma_V, \sigma_W) * E') \\
&= \sum_{(\sigma_V, \sigma_W) \in S_m \times S_n} \mu((\sigma_V, \sigma_W) * (X \setminus E')) \\
&= \sum_{(\sigma_V, \sigma_W) \in S_m \times S_n} \mu(X \setminus E') \\
&< \sum_{(\sigma_V, \sigma_W) \in S_m \times S_n} \frac{\epsilon}{|S_m \times S_n|} \\
&= \epsilon.
\end{aligned}$$

Note that the three conditions in Theorem E.1 are satisfied by the definition of Φ_{solu} , Corollary B.7, and Corollary B.6, respectively. Using Theorem E.1, there exists $F_W \in \mathcal{F}_{\text{GNN}}^W$ such that

$$\sup_{(G, H) \in E} \|F_W(G, H) - \Phi_{\text{solu}}(G, H)\| < \delta.$$

Therefore, it holds that

$$\text{Meas}(\{(G, H) \in X : \|F_W(G, H) - \Phi_{\text{solu}}(G, H)\| > \delta\}) \leq \text{Meas}(X \setminus E) < \epsilon,$$

which completes the proof. \square

Proof of Corollary 3.7. One can assume that \mathcal{D} is closed under the action of $S_m \times S_n$; otherwise, a larger but still finite dataset, $\bigcup_{(\sigma_V, \sigma_W) \in S_m \times S_n} (\sigma_V, \sigma_W) * \mathcal{D}$, can be considered instead of \mathcal{D} . The rest of the proof is similar to that of Theorem 3.6 since \mathcal{D} is compact and $\Phi_{\text{solu}}|_{\mathcal{D}}$ is continuous. \square

APPENDIX G. DETAILS OF THE NUMERICAL EXPERIMENTS

LP instance generation. We generate each LP with the following way. We set $m = 10$ and $n = 50$. Each matrix A is sparse with 100 nonzero elements whose positions are sampled uniformly and values are sampled normally. Each element in b, c are sampled i.i.d and uniformly from $[-1, 1]$. The variable bounds l, u are sampled with $\mathcal{N}(0, 10)$. If $l_j > u_j$, then we swap l_j and u_j for all $1 \leq j \leq n$. Furthermore, we sample \circ_i i.i.d with $\mathbb{P}(\circ_i = \text{"} \leq \text{"}) = 0.7$ and $\mathbb{P}(\circ_i = \text{"} = \text{"}) = 0.3$. With the generation approach above, the probability that each LP to be feasible is around 0.53.

MLP architectures. As we mentions in the main text, all the learnable functions in GNN are taken as MLPs. The input functions $f_{\text{in}}^V, f_{\text{in}}^W$ have one hidden layer and other functions $f_{\text{out}}, f_{\text{out}}^W, \{f_l^V, f_l^W, g_l^V, g_l^W\}_{l=0}^L$ have two hidden layers. The embedding size d_0, \dots, d_L are uniformly taken as d that is chosen from $\{2, 4, 8, 16, 32, 64, 128, 256, 512\}$. All the activation functions are ReLU.

Training settings. We use Adam [31] as our training optimizer with learning rate of 0.0003. The loss function is taken as mean squared error. All the experiments are conducted on a Linux server with an Intel Xeon Platinum 8163 GPU and eight NVIDIA Tesla V100 GPUs.

(Z. CHEN) DEPARTMENT OF MATHEMATICS, DUKE UNIVERSITY, DURHAM, NC 27708.

Email address: `ziang@math.duke.edu`

(J. LIU) DECISION INTELLIGENCE LAB, DAMO ACADEMY, ALIBABA US, BELLEVUE, WA 98004.

Email address: `jialin.liu@alibaba-inc.com`

(X. WANG) DECISION INTELLIGENCE LAB, DAMO ACADEMY, ALIBABA US, BELLEVUE, WA 98004.

Email address: `xinshang.w@alibaba-inc.com`

(J. LU) DEPARTMENTS OF MATHEMATICS, PHYSICS, AND CHEMISTRY, DUKE UNIVERSITY, DURHAM, NC 27708.

Email address: `jianfeng@math.duke.edu`

(W. YIN) DECISION INTELLIGENCE LAB, DAMO ACADEMY, ALIBABA US, BELLEVUE, WA 98004.

Email address: `wotao.yin@alibaba-inc.com`